

Relating subsets of a poset, and a partition theorem for WQOs

Reinhard Diestel

We point out some basic properties of the partial ordering which a poset P induces on its power set, defining $A \leq B$ to mean that every element of A lies below some element of B . One result is that if P is a WQO then P decomposes uniquely into finitely many indivisible sets A_1, \dots, A_n (that are essential parts of P in the sense that $P \not\leq P \setminus A_i$).

This note collects together some observations I made in the context of comparing graph properties by the graph minor relation [2], but which apply more generally to arbitrary subsets of a given poset. They are all simple and ought to be well known, but I have been unable to find a source. The main observation is that the subsets of an infinite poset P can be decomposed in a way that resembles factoring: there are ‘indivisible’ sets that behave like primes, and if P is a WQO then it factors uniquely into such indivisible sets.

Let (P, \leq) be any poset, typically infinite. (Countable will do to make things interesting, and a particularly interesting case will be that P is a WQO.) Given subsets $A, B \subseteq P$, let us write $A \leq B$ to express that for every $a \in A$ there is a $b \in B$ such that $a \leq b$. This is a quasi-ordering on the power set of P , which induces a partial ordering on the set \mathcal{P} of \sim -equivalence classes of subsets of P , where $A \sim B$ if $A \leq B$ and $B \leq A$. Although we shall continue to speak about the subsets themselves rather their equivalence classes, we shall often distinguish them only up to equivalence to derive properties of this poset \mathcal{P} . For $A, B \subseteq P$ we write $A < B$ if $A \leq B$ but $A \not\sim B$ (ie. $B \not\leq A$).

The first problem we address is whether we can always find particularly typical representatives of these equivalence classes, in the following sense. Given an infinite set $A \subseteq P$, we can obtain numerous equivalent sets just by ‘adding junk’: for every $A' < A$ we clearly have $A \cup A' \sim A$. This process is not easily reversible: if we are given $A \cup A'$ as a single set, we may not be able to identify and discard its ‘inessential’ part A' . So it seems that ‘lean’ sets not containing large amounts of such junk are particularly desirable representatives of their equivalence classes.

To make this precise, let us call an infinite set $A \subseteq P$ *lean* if $A \leq A'$ for every $A' \subseteq A$ with $|A'| = |A|$. (For example, the set of finite stars and the set of finite paths are both lean under the minor relation for graphs, but the set of finite trees is not lean.) Then our first question is: which infinite sets $A \subseteq P$ are equivalent to some lean set $A' \subseteq P$, possibly with $A' \subseteq A$?

Clearly some are not: the union U of the set of finite stars and the set of finite paths, for instance, is not equivalent to any lean set of finite graphs. The

reason is that U splits naturally into its two constituent subsets of stars and paths—which are incomparable under \leq and hence cannot be equivalent to all of U . One of our aims below will be to show that, for countable sets A , such splitting is the only possible obstruction to the existence of an equivalent lean set. So let us make this splitting precise.

Let A be any subset of P . We call $A' \subseteq A$ *small in A* if $A' < A$ (ie. if $A \not\leq A'$), and *large in A* if $A \leq A'$. The complements of small sets we call ‘essential’; thus, $A' \subseteq A$ is *essential in A* if $A \not\leq A \setminus A'$, or equivalently if $A' \not\leq A \setminus A'$. Note that subsets of a small subset of A are also small in A (and hence supersets of large or essential sets are again large or essential, respectively), and that an essential subset A' of A is also essential in every set A'' with $A' \subseteq A'' \subseteq A$.

If A is the union of two small subsets $A_1, A_2 \subseteq A$, we call A *divisible*. Replacing A_2 by $A_2 \setminus A_1$ if necessary we can always ensure that these small subsets are disjoint; we then say that A *splits into* these two subsets.

The small subsets of an indivisible set A form a set-theoretic ideal: their finite unions (as well as their subsets) are again small in A . This follows from the following factoring lemma, in which the indivisible sets appear as primes:

Lemma 1. *If A is indivisible and $A \leq B_1 \cup B_2$, then $A \leq B_1$ or $A \leq B_2$.*

Proof. For $i = 1, 2$ put $A_i := \{a \in A \mid \exists b \in B_i : a \leq b\}$. Our assumption of $A \leq B_1 \cup B_2$ implies that $A = A_1 \cup A_2$. So as A is indivisible the A_i cannot both be small, ie. one of them satisfies $A \leq A_i \leq B_i$. \square

Lemma 1 implies that, unlike leanness, divisibility and indivisibility are invariant under equivalence:

Corollary 2. *If A is indivisible and $A \sim B$ then B is indivisible.*

Proof. If $B = B_1 \cup B_2$ then $B \leq A \leq B_i$ for some $i \in \{1, 2\}$ by Lemma 1, so B_i is not small. \square

Since lean sets are indivisible, Corollary 2 implies that only indivisible sets can be equivalent to lean sets. But is every infinite indivisible set equivalent to some lean set, perhaps even a lean subset? In general, this looks like a difficult problem: the naive approach of recursively splitting small subsets off a given indivisible set ‘until it becomes lean’ is obviously fraught with problems; for example, it may happen that after any finite number of steps one still has a non-lean indivisible set but after ω steps the entire set has disappeared.

For countable sets, however, there is a very simple characterization of divisibility which essentially implies a positive answer to the above question:

Proposition 3. *For every countable set $A \subseteq P$ the following statements are equivalent:*

- (i) A is indivisible;
- (ii) A has an equivalent lean subset or a greatest element;
- (iii) A contains a chain C such that $A \leq C$;
- (iv) every two elements of A have a common upper bound in A .

Proof. (i) \rightarrow (iv) Suppose $a_1, a_2 \in A$ have no common upper bound in A . Then the sets $A_1 := \{a \in A \mid a_1 \not\leq a\}$ and $A_2 := \{a \in A \mid a_2 \not\leq a\}$ are small in A but have union A , so A is divisible.

(iv) \rightarrow (iii) Enumerate A as $\{a_1, a_2, \dots\}$ and use (iv) to construct the desired chain $c_1 \leq c_2 \leq \dots$ inductively, choosing each c_n above (\geq) both a_n and c_{n-1} .

(iii) \rightarrow (ii) If C has a maximal element, then this is the greatest element of A . If not, then C has a cofinal subchain of order type ω . This subchain is lean and equivalent to C , and hence also to A .

(ii) \rightarrow (i) If A has a greatest element x then a subset of A is large if and only if it contains x . So A cannot split into two small subsets (which would also be large). If A is equivalent to a lean (and thus indivisible) set, then A is indivisible by Corollary 2. \square

What about uncountable sets A ? As it turns out, the equivalence of (i) and (iv) above remains valid, but these conditions no longer imply (ii) and (iii):

Proposition 4. *An arbitrary set $A \subseteq P$ is indivisible if and only if every two elements of A have a common upper bound in A .*

Proof. The forward implication was shown in the proof of Propositions 3, which made no use of countability in this part. For the converse, let $A = A_1 \cup A_2$ and assume that A_1 is small; we show that A_2 is large, ie. that $A \leq A_2$.

Since $A \not\leq A_1$, there exists an $x \in A$ such that A_1 fails to meet the up-closure $[x] = \{a \in A \mid x \leq a\}$ of x in A . So $[x] \subseteq A_2$. By assumption, there exists for every $a \in A$ some $b \in A$ such that $a, x \leq b$, ie. with $b \in [x] \subseteq A_2$. So $A \leq A_2$ as desired. \square

Here is an example of an uncountable poset P that is indivisible but fails (for $A = P$) to satisfy (ii) and (iii) of Proposition 3. We construct P recursively in ω steps, starting with a set of \aleph_1 minimal elements. At each step, we add one element z above every pair x, y of elements from previous steps, leaving z unrelated to all the other elements.

By construction, any two elements of P have a common upper bound in P , so by Proposition 4 the set P is indivisible. On the other hand, every down-set $[x]$ is finite, and so any chain C in P (which is clearly countable) has a countable down-closure and thus fails to satisfy $P \leq C$ as required in (iii). For

the same reason, any $A \subseteq P$ equivalent to P must be uncountable. Then A meets one of the ω levels of P in an uncountable subset A' . Thus $|A'| = |A| = \aleph_1$ but $P \not\leq A'$, and hence $A \not\leq A'$ as $A \sim P$. Therefore A is not lean, showing that P is a counterexample to (ii).

Let us now leave the subject of equivalent lean sets or subsets, and take a closer look at divisible and indivisible sets A for their own sake. Since now all that matters is the structure of A , we lose nothing by just considering the divisibility of P itself.

Lemma 1 suggests that the analogy to factoring which the word ‘divisibility’ suggests may be deeper than one would at first expect. And indeed, we have the following pretty ‘prime factor theorem’ for WQOs:

Theorem 5. *If P is a WQO, then P can be partitioned into finitely many indivisible essential subsets A_1, \dots, A_n . This partition is unique up to equivalence; in fact, every indivisible essential subset of P is equivalent to one of the A_i .*

For the proof we need as a lemma the forward implication of the following observation relating the ordering on P to that on its power set. Again, I expect this lemma to be folklore among WQO specialists but include the proof for completeness.

Lemma 6. *P is a WQO if and only if all chains $A_1 > A_2 > \dots$ of subsets of P are finite.*

Proof. For the forward implication, let $A_1 > A_2 > \dots$ be a strictly descending chain of subsets of P . Since $A_1 \not\leq A_2$, there is an $a_1 \in A_1$ such that $a_1 \not\leq a$ for all $a \in A_2$. Then $a_1 \not\leq a$ even for all $a \in A_i$ with $i \geq 2$, because $A_i \leq A_2$. Similarly, A_2 has an element $a_2 \not\leq a$ for all $a \in A_i$ with $i \geq 3$, and so on. We thus obtain a sequence a_1, a_2, \dots with $a_i \in A_i$ for each i and such that $a_i \not\leq a_j$ whenever $i < j$. As P is a WQO every such sequence is finite, and hence so is our chain $A_1 > A_2 > \dots$.

Conversely, if P is not a WQO it contains a bad sequence a_1, a_2, \dots : an infinite sequence such that $a_i \not\leq a_j$ for all $i < j$. Then the tails $\{a_i, a_{i+1}, \dots\}$ of this sequence form an infinite strictly descending chain of subsets of P . \square

Proof of Theorem 5. Let us begin by constructing some sets A_1^*, \dots, A_n^* similar to the desired partition sets A_1, \dots, A_n . We do this recursively in finitely many steps, at each step modifying a finite list \mathcal{L} of subsets of P until no further modifications are possible. We start with the one-element list $\mathcal{L} = \{P\}$. At each step of the construction, we first check whether any element of \mathcal{L} is inessential in $\bigcup \mathcal{L}$; if so, we delete it from the list and proceed to the next step. If not, we try to find a divisible element of \mathcal{L} . If none exists, we terminate the construction. If some $A \in \mathcal{L}$ is divisible, we split it into disjoint small subsets A' and A'' to replace A in \mathcal{L} , and proceed to the next step.

If this construction continues indefinitely, then by König's infinity lemma [1] there is an infinite chain $C_1 > C_2 > \dots$ of subsets of P (each chosen as a small subset of the previous), which by Lemma 6 cannot exist. So the construction terminates after finitely many steps, with a list $\mathcal{L}^* = \{A_1^*, \dots, A_n^*\}$ of disjoint indivisible subsets of P that are essential in $P^* := \bigcup \mathcal{L}^*$. Since every construction step (in particular, the deletion of inessential subsets) preserves the initial property of \mathcal{L} that $P \leq \bigcup \mathcal{L}$, we have $P \leq P^*$. We can therefore find a partition $A_1 \cup \dots \cup A_n$ of P such that

$$A_i \leq A_i^* \subseteq A_i \tag{1}$$

(and hence $A_i \sim A_i^*$) for every i : the sets

$$A'_i := \{x \in P \mid \exists a \in A_i^*: x \leq a\} \supseteq A_i^*$$

cover P , and to obtain the A_i we just make them disjoint by trimming off any overlaps outside the (disjoint) A_i^* ; then $A_i^* \subseteq A_i \subseteq A'_i \leq A_i^*$ for all i , as required.

Since the A_i^* are indivisible and $A_i \sim A_i^*$, the A_i are indivisible by Corollary 2. Let us now show that every A_i is essential in P . If not, then $A_i \leq \bigcup_{j \neq i} A_j$ and therefore

$$A_i^* \subseteq A_i \leq \bigcup_{j \neq i} A_j \leq \bigcup_{j \neq i} A_j^*$$

by (1), which contradicts the fact that A_i^* is essential in P^* .

To complete the proof, it remains to show that every essential indivisible subset B of P is equivalent to one of the sets A_1, \dots, A_n . For each $i = 1, \dots, n$, put $B_i := B \cap A_i$. Then $B = B_1 \cup \dots \cup B_n$, and since B is indivisible we have $B \leq B_i$ for some i by Lemma 1, and hence $B \sim B_i \subseteq A_i$. If $B \sim A_i$ we are done; if not, then B_i is small in A_i . Then $A_i \setminus B_i$ is large in A_i (because A_i is indivisible), ie. $A_i \leq A_i \setminus B_i$. Thus

$$B \leq B_i \subseteq A_i \leq A_i \setminus B_i \subseteq P \setminus B,$$

which contradicts our assumption that B is essential in P . □

We remark that the essential sets A_1^*, \dots, A_n^* from the proof of Theorem 5 can easily be 'constructed' directly (and seen to be essential in P), as follows. Note first that a subset of P is essential if and only if it contains the entire up-closure $\lfloor x \rfloor$ in P of one of its points x . Using Proposition 4 and the fact that P contains no infinite antichain, one easily shows that above each point of P there is a point x whose up-closure $\lfloor x \rfloor$ in P is indivisible. Thus if $\{a_1, \dots, a_n\}$ is any maximal set of points with disjoint indivisible up-closures, then these up-closures $A_i^* := \lfloor a_i \rfloor$ are indivisible essential subsets of P such that

$P \leq A_1^* \cup \dots \cup A_n^*$. The partition $P = A_1 \cup \dots \cup A_n$ for the theorem is then obtained exactly as in the proof: by replacing each A_i^* with its down-closure and making these disjoint by trimming off overlaps.

Note that in this proof we only use that P has no infinite antichain, or indeed that P contains no induced copy of the binary tree. Therefore either of these assumptions can replace the stronger WQO assumption in Theorem 5. A full characterization of the countable posets that factor into indivisible essential subsets will be given in [3]; the corresponding problem for larger posets remains open.

References

- [1] R. Diestel, *Graph theory*, 2nd edition, Springer-Verlag 2000 and <http://www.math.uni-hamburg.de/home/diestel/books/graph.theory/download.html>
- [2] R. Diestel & D. Kühn, Graph minor hierarchies, preprint 2001.
- [3] R. Diestel & O. Pikhurko, On the cofinality of infinite partially ordered sets: factoring a poset into lean essential subsets, in preparation.