

Exercises in Algebra (master): Homological Algebra

Prof. Dr. Birgit Richter

Summer term 2018

Exercise sheet no 5

for the exercise class on the 16th of May 2018

1 (Suspensions)

Let $p \in \mathbb{Z}$ be arbitrary and let (C_*, d_C) be a chain complex. Define $\Sigma^p C_*$ as

$$(\Sigma^p C)_n = C_{n-p} \text{ and } d_{\Sigma^p C}(c) = (-1)^p d_C(c).$$

Calculate $H_n \Sigma^p C_*$ in terms of the homology groups of C_* .

2 (Mapping cones)

Let $f_*: (C_*, d_*) \rightarrow (C'_*, d'_*)$ be a chain map. We define the *mapping cone of f* as $C(f)_n = (C_{n-1}, C'_n)$ and we let $\tilde{d}: C(f)_n \rightarrow C(f)_{n-1}$ be the map that sends (c, c') to $(-d_{n-1}(c), d'_n(c') - f_{n-1}(c))$.

- (1) Check that this actually defines a chain complex.
- (2) Prove that there is a short exact sequence of chain complexes

$$0 \longrightarrow C'_* \xrightarrow{j} C(f)_* \xrightarrow{e} \Sigma C_* \longrightarrow 0.$$

What is the connecting homomorphism in this case?

- (3) Show that a chain map $g_*: C_* \rightarrow D_*$ is chain homotopic to the zero map if and only if g_* extends over the mapping cone of the identity map of C_* :

$$\begin{array}{ccc} C_* & \xrightarrow{g_*} & D_* \\ \downarrow j & \nearrow & \\ C(\text{id}_{C_*}) & & \end{array}$$

3 (Simple but important chain complexes)

- (1) Let A be any abelian group and let n be an integer. We consider the chain complex $S^n(A)$ that has A in degree n and is trivial in all other degrees. What is $H_m(S^n(A))$ for all m ? What is the abelian group of chain maps from $S^n(\mathbb{Z})$ to any chain complex C_* ?
- (2) Let A and n be as above and let $D^n(A)$ be the chain complex with $D^n(A)_n = A = D^n(A)_{n-1}$ and $D^n(A)_k = 0$ for all $k \notin \{n, n-1\}$. As the differential we take the identity map of A between chain degrees n and $n-1$ and take the zero map everywhere else. What is $H_m(D^n(A))$ for all m ? What is the abelian group of chain maps from $D^n(\mathbb{Z})$ to any chain complex C_* ?
- (3) Let $f \in \text{Ab}(A, B)$. Consider f as a chain complex by placing A in degree one, B in degree zero and using f as the only non-trivial boundary operator. What are the homology groups of this chain complex?