

Exercises in Algebraic Topology (master)

Prof. Dr. Birgit Richter

Summer term 2017

Exercise sheet no 0

due: 10th of April 2017

As we have no-one who will correct the exercises, 'due' means that you should work on the exercises and should be able to present solutions during the exercise class on the 10th of April.

1 (Abelian groups) Let A and B be two abelian groups. We denote by $\text{Hom}(A, B)$ the set of group homomorphisms from A to B .

a) Show that $\text{Hom}(A, B)$ is an abelian group.

b) What is $\text{Hom}(\mathbb{Z}, A)$?

c) Let $n > 1$ be a natural number. Describe $\text{Hom}(\mathbb{Z}/n\mathbb{Z}, A)$ as a subgroup of A . What is $\text{Hom}(\mathbb{Z}/n\mathbb{Z}, \mathbb{Z})$ or $\text{Hom}(\mathbb{Z}/n\mathbb{Z}, \mathbb{Q})$?

2 (Disks and spheres)

Let \mathbb{D}^n be the chain complex whose only non-trivial entries are in degrees n and $n-1$ with $\mathbb{D}_n^n = \mathbb{D}_{n-1}^{n-1} = \mathbb{Z}$. Its only non-trivial boundary operator is the identity.

Similarly, let \mathbb{S}^n be the chain complex whose only non-trivial entry is in degree n with $\mathbb{S}_n^n = \mathbb{Z}$.

a) Assume that (C_*, d) is an arbitrary chain complex. Describe chain maps from \mathbb{D}^n to C_* and from \mathbb{S}^n to C_* in terms of subobjects of C_* .

b) Are there chain maps between \mathbb{D}^n and \mathbb{S}^m ? What is the homology of \mathbb{D}^n and \mathbb{S}^m ?

c) Let $f_*: C_* \rightarrow C'_*$ be a chain map and assume that f_n is a monomorphism for all n . Do we then know that $H_n(f_*)$ is also a monomorphism?

3 (Too much to ask for?)

a) What are the homology groups of the chain complex

$$C_* = (\dots \rightarrow \mathbb{Z}/4\mathbb{Z} \xrightarrow{2\cdot} \mathbb{Z}/4\mathbb{Z} \xrightarrow{2\cdot} \mathbb{Z}/4\mathbb{Z} \rightarrow \dots)?$$

b) Is there a chain homotopy from the identity of C_* to the zero map, *i.e.*, are there maps $s_n: C_n \rightarrow C_{n+1}$ with $d \circ s + s \circ d = \text{id}_{C_n}$ for all $n \in \mathbb{Z}$?

4 (Lego)

Let $(A_n)_{n \in \mathbb{Z}}$ be an arbitrary family of finitely generated abelian groups. Is there a chain complex F_* with F_n free abelian for all $n \in \mathbb{Z}$ and with $H_n(F_*) \cong A_n$? (Recall the structure theorem for finitely generated abelian groups for this.)