

# The inner automorphism 3-group of a strict 2-group

David Roberts and Urs Schreiber

August 13, 2007

## Abstract

Any group  $G$  gives rise to a 2-group of inner automorphisms,  $\text{INN}(G)$ . It is an old result by Segal that the nerve of this is the universal  $G$ -bundle. We discuss that, similarly, for every 2-group  $G_{(2)}$  there is a 3-group  $\text{INN}(G_{(2)})$  and a slightly smaller 3-group  $\text{INN}_0(G_{(2)})$  of inner automorphisms. We describe these for  $G_{(2)}$  any strict 2-group, discuss how  $\text{INN}_0(G_{(2)})$  can be understood as arising from the mapping cone of the identity on  $G_{(2)}$  and show that its underlying 2-groupoid structure fits into a short exact sequence

$$G_{(2)} \longrightarrow \text{INN}_0(G_{(2)}) \longrightarrow \Sigma G_{(2)} .$$

As a consequence,  $\text{INN}_0(G_{(2)})$  encodes the properties of the universal  $G_{(2)}$  2-bundle.

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b><math>n</math>-Groups in terms of groups</b>	<b>7</b>
2.1	Conventions for strict 2-groups and crossed modules . . . . .	7
2.2	3-Groups and 2-crossed modules . . . . .	10
2.3	Mapping cones of crossed modules . . . . .	12
<b>3</b>	<b>Main results</b>	<b>13</b>
3.1	The exact sequence $G_{(2)} \rightarrow \text{INN}_0(G_{(2)}) \rightarrow \Sigma G_{(2)}$ . . . . .	13
3.2	$\text{INN}_0(G_{(2)})$ from a mapping cone . . . . .	13
<b>4</b>	<b>Inner automorphism <math>n</math>-groups</b>	<b>13</b>
4.1	Exact sequences of strict 2-groupoids . . . . .	14
4.2	Tangent 2-categories . . . . .	15
4.3	Inner automorphisms . . . . .	18
4.4	Inner automorphism 3-groups. . . . .	20

<b>5</b>	<b>The 3-group <math>\text{INN}_0(G_{(2)})</math></b>	<b>24</b>
5.1	Objects . . . . .	24
5.2	Morphisms . . . . .	24
5.2.1	Composition . . . . .	25
5.2.2	Product . . . . .	25
5.3	2-Morphisms . . . . .	28
5.3.1	Composition . . . . .	28
5.3.2	Product . . . . .	31
<b>6</b>	<b>Properties of <math>\text{INN}_0(G_{(2)})</math></b>	<b>32</b>
6.1	Structure morphisms . . . . .	32
6.1.1	Strictness as a 2-groupoid . . . . .	32
6.2	Trivializability . . . . .	33
6.3	Universality . . . . .	34
6.4	The corresponding 2-crossed module . . . . .	34
6.4.1	Relation to the mapping cone of $H \rightarrow G$ . . . . .	41
<b>7</b>	<b>Universal <math>n</math>-bundles</b>	<b>42</b>
7.1	Universal 1-bundles in terms of $\text{INN}(G)$ . . . . .	42
7.2	Universal 2-bundles in terms of $\text{INN}_0(G_{(2)})$ . . . . .	44

## 1 Introduction

The theory of groups and their principal fiber bundles generalizes to that of categorical groups and their categorical principal fiber bundles. In fact, using higher categories, one has for each integer  $n$  the notion of  $n$ -groups and their principal  $n$ -bundles.

The reader may have encountered principal 2-bundles mostly in the language of (nonabelian) gerbes, which are to 2-bundles essentially like sheaves are to ordinary bundles. The concept of a 2-bundle proper is described in [12, 13].

These  $n$ -bundles are certainly interesting already in their own right. One crucial motivation for considering them comes from the study of  $n$ -dimensional quantum field theory. In this case one is interested in  $n$ -dimensional analogs of the concept of parallel transport in fiber bundles with connection [15, 16].

In that context a curious phenomenon occurs: whenever one investigates  $n$ -dimensional quantum field theory governed by an  $n$ -group  $G_{(n)}$ , it turns out [8] that the situation is governed by an  $(n+1)$ -group associated to  $G_{(n)}$ . In fact, it is appropriate to call this  $(n+1)$ -group  $\text{INN}_0(G_{(n)})$ , because, as the notation suggests, it is related to inner automorphisms of the original  $n$ -group  $G_{(n)}$ .

One of our aims here is to define what inner automorphisms of a 2-group are and to give a concise definition as well as a detailed description of the 3-group  $\text{INN}_0(G_{(2)})$  for any strict 2-group  $G_{(2)}$ . We then prove that  $\text{INN}_0(G_{(2)})$  has a couple of rather peculiar properties; it is *trivializable* (equivalent, as a 2-group, to the trivial 2-group), and fits into a short exact sequence

$$G_{(2)} \hookrightarrow \text{INN}_0(G_{(2)}) \twoheadrightarrow \Sigma G_{(2)}$$

of 2-groupoids. To appreciate this result, it is helpful to first consider the analogous statement for ordinary groups.

**The statement for ordinary groups.** For any ordinary group  $G$ , various constructions of interest, like that of the universal  $G$ -bundle, are closely related to a certain groupoid determined by  $G$ .

There are several different ways to think of this groupoid. The simplest way to describe its structure is to say that it is the codiscrete groupoid over the elements of  $G$ , namely the groupoid whose objects are the elements of  $G$  and which has precisely one morphism from any element to any other.

The relevance of this groupoid is better understood by thinking of it as the action groupoid  $G//G$  of the action of  $G$  on itself by left multiplication. As such, we may write any of its morphisms as

$$g \xrightarrow{h} hg$$

for  $g, h \in G$  and  $hg$  being the product of  $h$  and  $g$  in  $G$ .

While this way of thinking about our groupoid already makes it more plausible that it is related to  $G$ -actions and hence possibly to  $G$ -bundles, one more property remains to be made manifest: there is also a monoidal structure on our groupoid. For any two morphisms,

$$g_1 \xrightarrow{h_1} h_1 g_1$$

and

$$g_2 \xrightarrow{h_2} h_2 g_2,$$

we can form the product morphism

$$g_2 g_1 \xrightarrow{h_2 \text{Ad}_{g_2}(h_1)} h_2 g_2 h_1 g_1,$$

and this assignment is functorial in both arguments. Moreover, to every morphism

$$g \xrightarrow{h} hg$$

there is a morphism

$$g^{-1} \xrightarrow{\text{Ad}_{g^{-1}}(h)^{-1}} (hg)^{-1},$$

which is its inverse with respect to this product operation.

This makes  $G//G$  a strict 2-group [1]. A helpful way to make the 2-group structure on  $G//G$  more manifest is to relate it to inner automorphisms of  $G$ . To see this, consider another groupoid canonically associated to any group  $G$ , namely the groupoid

$$\Sigma G = \{ \bullet \xrightarrow{g} \bullet \mid g \in G \}$$

which has a single object  $\bullet$ , one morphism for each element of  $G$  and where composition of morphisms is just the product in the group.

Automorphisms

$$a : \Sigma G \rightarrow \Sigma G$$

(i.e. invertible functors) of this groupoid are nothing but group automorphisms of  $G$ . But now there are also isomorphisms between two such morphisms  $a$  and  $a'$ , namely natural transformations:

$$\begin{array}{ccc} & a & \\ \Sigma G & \begin{array}{c} \Downarrow \\ \Downarrow \end{array} & \Sigma G \\ & a' & \end{array} .$$

This way for every ordinary group  $G$  we have not just its ordinary group of automorphisms, but actually a 2-group

$$\text{AUT}(G) := \text{Aut}_{\text{Cat}}(\Sigma G) .$$

This is a groupoid, whose objects are group automorphisms of  $G$ . The 2-group structure on this groupoid is manifest from the horizontal composition of the natural transformations above. Hence the ordinary automorphism group of  $G$  is the group of objects of  $\text{AUT}(G)$ .

By writing out the definition of a natural transformation, one sees that there is a morphism between two objects in  $\text{AUT}(G)$  whenever the two underlying ordinary automorphisms of  $G$  differ by conjugation with an element of  $G$ . It follows in particular that the *inner* automorphisms of  $G$  correspond to those autofunctors of  $\Sigma G$  which are isomorphic to the identity:

$$\begin{array}{ccc} & \text{Id} & \\ \Sigma G & \begin{array}{c} \Downarrow g \simeq \\ \Downarrow \end{array} & \Sigma G \\ & \text{Ad}_g & \end{array} .$$

Therefore consider the groupoid  $\text{INN}(G)$ : its objects are pairs, consisting of an automorphisms together with a transformation connecting it to the identity. A morphism from  $(g, \text{Ad}_g)$  to  $(gh, \text{Ad}_{gh})$  is a commuting triangle

$$\begin{array}{ccc} & g & \rightarrow \text{Ad}_g \\ \text{Id}_{\Sigma G} & \begin{array}{c} \Downarrow \\ \Downarrow \end{array} & \downarrow h \\ & hg & \rightarrow \text{Ad}_{hg} \end{array} .$$

This is again exactly the groupoid  $G//G$  which we are discussing

$$\text{INN}(G) = G//G.$$

In this formulation the natural notion of composition of group automorphisms nicely explains the monoidal structure on  $G//G$ .

Notice that  $\text{INN}(G)$  remembers the center of the group. We will discuss that it sits inside an exact sequence

$$1 \rightarrow Z(G) \rightarrow \text{INN}(G) \rightarrow \text{AUT}(G) \rightarrow \text{OUT}(G) \rightarrow 1$$

of 2-groups, and that this is what generalizes to higher  $n$ .

If we think of the group  $G$  just as a discrete category, whose objects are the elements of  $G$  and which has only identity morphisms, then there is an obvious monomorphic functor

$$G \rightarrow \text{INN}(G).$$

Moreover, there is an obvious epimorphic functor

$$\text{INN}(G) \rightarrow \Sigma G$$

from our groupoid to the group  $G$ , but now with the latter regarded as a category with a single object. This simply forgets the source and target labels and recalls only the group element which is acting.

These two functors are such that the image of the former is precisely the collection of morphisms which get sent to the identity morphism by the latter. Therefore we say that we have a short exact sequence

$$G \rightarrow \text{INN}(G) \rightarrow \Sigma G \tag{1}$$

of groupoids.

Notice that  $G$  and  $\text{INN}(G)$  are groupoids which are also 2-groups (the first one, being an ordinary group, is a degenerate case of a 2-group), and that the morphism  $G \rightarrow \text{INN}(G)$  is also a morphism of 2-groupoids. But  $\Sigma G$  is in general just a groupoid without monoidal structure – it has the structure of a 2-group if and only if  $G$  is abelian.

Even though all this is rather elementary, the exact sequence (1) is important. We can apply the functor  $|\cdot|$  to our sequence, which takes the nerve of a category and then forms the geometric realization. Note that when  $K$  is a 2-group,  $|K|$  is a topological group. Under this functor, (1) becomes the universal  $G$ -bundle

$$G \rightarrow EG \rightarrow BG,$$

even when  $G$  is a topological or Lie group. The fact that  $BG \simeq |\Sigma G|$  is the very definition of the classifying space  $BG$  of a group  $G$  [2]. That  $EG \simeq |\text{INN}(G)|$  is contractible follows from the existence of an equivalence of groupoids  $\text{INN}(G) \xrightarrow{\sim} *$ . Finally, the inclusion  $G \rightarrow \text{INN}(G)$  together with the monoidal structure on  $\text{INN}(G)$  gives the free  $G$ -action of  $G$  on  $EG$  whose quotient is exactly  $BG$ . The observation that  $|\text{INN}(G)|$  is a model for  $EG$  is originally due to Segal [2].

Our first main result is the higher categorical analogue of (1), obtained by starting with a strict 2-group  $G_{(2)}$  in place of the ordinary group  $G$ .

Here we do not consider geometric realizations of our categories and 2-categories (for more on that see the closely related article [17]) but instead focus on the existence of these sequences of 1- and 2-groupoids. We comment on further aspects of the topic of universal  $n$ -bundles in §7. More details will be given in [14].

**The formulation in terms of crossed modules.** For many purposes, like doing explicit computations and for applying the rich toolbox of simplicial methods, it is possible (and useful!) to express  $n$ -groups in terms of  $n$ -term complexes of ordinary groups with extra structure on them. For instance strict 2-groups are well known to be equivalent to crossed modules of two ordinary groups: one describes the group of objects, the other the group of morphisms of the 2-group.

This pattern continues, but there is a bifurcation of constructions, all of which are equivalent. Sufficiently strict 3-groups are described by 2-crossed modules, which involve three ordinary groups forming a normal complex, and also by crossed squares, which look like crossed modules of crossed modules of groups. We will primarily use the former, and only mention crossed squares when we cannot avoid it.<sup>1</sup> The way we use these two models can be illustrated in one lower categorical dimension by comparing the *map*

$$G \xrightarrow{\text{id}} G$$

to the *crossed module*

$$G \xrightarrow{\text{id}} G \xrightarrow{\text{Ad}} \text{Aut}(G)$$

using that map. The crossed module can be thought of as the mapping cone (=homotopy quotient) of the identity map.

The translation between  $n$ -groups and their corresponding  $n$ -term complexes of ordinary groups sheds light on both of these points of view. The analogue of our statement about the 3-group  $\text{INN}_0(G_{(2)})$  is our second main result: the complex of groups describing  $\text{INN}_0(G_{(2)})$  is the mapping cone of the identity on the complex of groups describing  $G_{(2)}$  itself.

This fact was anticipated from considerations in the theory of Lie  $n$ -algebras [8], where the Lie  $(n+1)$ -algebra corresponding to a Lie  $(n+1)$ -group  $\text{INN}_0(G_{(n)})$  has proven to be crucial for understanding connections with values in Lie  $n$ -algebras. There one finds that inner derivation Lie  $(n+1)$ -algebras govern Lie  $(n+1)$ -algebras of Chern-Simons type [8]. The fact that  $\text{INN}_0(G_{(2)})$  arises from a mapping cone of the identity is crucial in this context.

---

<sup>1</sup>As one goes to higher categorical dimensions (which we do not do here), there are multiple directions in which to extend the relevant diagrams, so there are a number of different models for  $n$ -groups. There is a sort of nonabelian Dold-Kan theorem, due to Carrasco and Cegarra [3], which can be used to characterise  $n$ -groups by  $n$ -term complexes of (possibly nonabelian) groups with the structure they call a hypercrossed complex.

**The plan of this article.** The main content of this work is as follows – first a concise and natural definition of inner automorphisms of 2-groups, relating them to the full automorphism  $(n+1)$ -group and to the categorical center. Then we apply this definition to the case that  $G_{(2)}$  is any strict 2-group and work out in full detail how  $\text{INN}_0(G_{(2)})$  looks like, i.e. how the various composition operations work, thus extracting its description in terms of complexes of ordinary groups. We state and prove the main properties of  $\text{INN}_0(G_{(2)})$ .

The plan of our discussion is as follows.

- In part 2 we recall the relation between 2- and 3-groups and crossed modules of ordinary groups. This serves to set up our convention for the precise choice of identification of 2-group morphisms with ordinary group elements.
- In part 3 we state our two main results.
- In part 4 we define inner automorphism  $n$ -groups and prove some important general properties of them.
- In part 5 we apply our definition of inner automorphisms to an arbitrary strict 2-group  $G_{(2)}$ , to form the 3-group  $\text{INN}_0(G_{(2)})$ . We then work out in detail the description of  $\text{INN}_0(G_{(2)})$  in terms of ordinary groups, spelling out the nature of the various composition and product operations.
- In part 6 we state and prove the main properties of  $\text{INN}_0(G_{(2)})$ , including our two main results.
- In part 7 we close by indicating in more detail how inner automorphism  $(n+1)$ -groups play the role of universal  $n$ -bundles.

We are grateful to Jim Stasheff for helpful discussions and for emphasizing the importance of the mapping cone construction in the present context. We profited from general discussion with Danny Stevenson and thank him for his help on the references to Segal's work. We also thank Christoph Schweigert and Zoran Škoda for helpful comments on the manuscript and Tim Porter for reminding us of Norrie's work.

## 2 $n$ -Groups in terms of groups

Sufficiently strict  $n$ -groups are equivalent to certain structures – crossed modules and generalizations thereof – involving just collections of ordinary groups with certain structure on them.

### 2.1 Conventions for strict 2-groups and crossed modules

An ordinary group  $G$  may be regarded as a one object category. If we regard  $G$  as such a category, we write  $\Sigma G$  in order to emphasize that we are thinking of the monoidal 0-category  $G$  as a one object 1-category.

This way we obtain a notion of  $n$ -groups from any notion of  $n$ -categories: an  $n$ -group  $G_{(n)}$  is a monoidal  $(n - 1)$ -category such that when regarded as a one-object  $(n)$ -category  $\Sigma G_{(n)}$  it becomes a one-object  $n$ -groupoid. An  $n$ -groupoid is an  $n$ -category all whose  $k$ -morphisms are equivalences, for all  $1 \leq k \leq n$ .

Here we shall be concerned with strict 2-groups and with 3-groups which are Gray categories. A strict 2-group  $G_{(2)}$  is one such that  $\Sigma G_{(2)}$  is a strict one-object 2-groupoid. A Gray groupoid is a 3-groupoid which is strict except for the exchange law of 2-morphisms.

The standard reference for 2-groups is [1]

It is well known that strict 2-groups are equivalent to crossed modules of ordinary groups.

**Definition 1.** A crossed module of groups is a diagram

$$H \xrightarrow{t} G \xrightarrow{\alpha} \text{Aut}(H)$$

in Grp such that

$$\begin{array}{ccc} H & \xrightarrow{\text{Ad}} & \text{Aut}(H) \\ & \searrow t & \nearrow \alpha \\ & G & \end{array}$$

and

$$\begin{array}{ccc} G \times H & \xrightarrow{\text{Id} \times t} & G \times G \\ \alpha \downarrow & & \downarrow \text{Ad} \\ H & \xrightarrow{t} & G \end{array} .$$

**Definition 2.** A strict 2-group  $G_{(2)}$  is any of the following equivalent entities

- a group object in Cat
- a category object in Grp
- a strict 2-groupoid with a single object

A detailed discussion can be found in [1].

One identifies

- $G$  is the group of objects of  $G_{(2)}$ .
- $H$  is the group of morphism of  $G_{(2)}$  starting at the identity object.
- $t : H \rightarrow G$  is the target homomorphism such that  $h : \text{Id} \rightarrow t(h)$  for all  $h \in H$ .



- $\alpha : G \rightarrow \text{Aut}(H)$  is conjugation with identity morphisms:

$$\text{Ad}_{\text{Id}_g}(\text{Id} \xrightarrow{h} t(h)) = \text{Id} \xrightarrow{\alpha(h)(h)} t(\alpha(g)(h))$$

for all  $g \in G, h \in H$ .

We often abbreviate

$$g_h := \alpha(g)(h).$$

Beyond that there are  $2 \times 2$  choices to be made when identifying a strict 2-group  $G_{(2)}$  with a crossed module of groups.

The first choice to be made is in which order to multiply elements in  $G$ . For

- $\bullet \xrightarrow{g_1} \bullet$  and  $\bullet \xrightarrow{g_2} \bullet$  two morphisms in  $\Sigma G_{(2)}$ , we can either set

$$\bullet \xrightarrow{g_1} \bullet \xrightarrow{g_2} \bullet := \bullet \xrightarrow{g_1 g_2} \bullet \quad (\text{F})$$

or

$$\bullet \xrightarrow{g_1} \bullet \xrightarrow{g_2} \bullet := \bullet \xrightarrow{g_2 g_1} \bullet \quad (\text{B}).$$

The other choice to be made is how to describe arbitrary morphisms by an element in the semidirect product group  $G \ltimes H$ : every morphism of  $G_{(2)}$  may be written as the product of one starting at the identity object with an identity morphism on some object. The choice of ordering here yields either

$$\begin{array}{c} g \\ \curvearrowright \\ \bullet \quad \parallel \quad \bullet \\ \downarrow h \\ \bullet \quad \curvearrowleft \\ \end{array} := \begin{array}{c} \text{Id} \\ \curvearrowright \\ \bullet \quad \parallel \quad \bullet \\ \downarrow h \\ \bullet \quad \curvearrowleft \\ \end{array} \xrightarrow{g} \bullet \quad (\text{R})$$

or

$$\begin{array}{c} g \\ \curvearrowright \\ \bullet \quad \parallel \quad \bullet \\ \downarrow h \\ \bullet \quad \curvearrowleft \\ \end{array} := \bullet \xrightarrow{g} \begin{array}{c} \text{Id} \\ \curvearrowright \\ \bullet \quad \parallel \quad \bullet \\ \downarrow h \\ \bullet \quad \curvearrowleft \\ \end{array} \quad (\text{L})$$

Here we choose the convention

$$LB.$$

This implies

$$\begin{array}{c} g \\ \curvearrowright \\ \bullet \quad \parallel \quad \bullet \\ \downarrow h \\ \bullet \quad \curvearrowleft \\ g' = t(h)g \\ \end{array}$$

for all  $g \in G$ ,  $h \in H$ , as well as the following two equations for horizontal and vertical composition in  $\Sigma G_{(2)}$ , expressed in terms of operations in the crossed module

and

## 2.2 3-Groups and 2-crossed modules

As we are considering strict models in this paper, we will assume that all 3-groups are as strict as possible. This means they will be one-object Gray-categories, or *Gray-monoids* [11]. A Gray-monoid is a (strict) 2-category  $\mathcal{M}$  such that the product functor

$$\mathcal{M} \otimes \mathcal{M} \rightarrow \mathcal{M}$$

uses the Gray tensor product, not the usual Cartesian product of 2-categories. Thus non-identity coherence morphisms only appear when we use the monoidal structure on  $\mathcal{M}$ .

Just as a 2-group gives rise to a crossed module, a 3-group gives rise to a 2-crossed module. Roughly, this is a complex of groups

$$L \rightarrow M \rightarrow N,$$

and a function

$$M \times M \rightarrow L \tag{2}$$

such that  $L \rightarrow M$  is a crossed module, and (2) measures the failure of  $M \rightarrow N$  to be a crossed module. An example is when  $L = 1$ , and then we have a crossed module. Now for the formal definition. See [5].

**Definition 3.** A **2-crossed module** is a normal complex of length 2

$$L \xrightarrow{\partial_2} M \xrightarrow{\partial_1} N$$

of  $N$ -groups ( $N$  acting on itself by conjugation) and an  $N$ -equivariant function

$$\{\cdot, \cdot\} : M \times M \rightarrow L,$$

called a *Peiffer lifting*, satisfying these conditions:

1.  $\partial_2\{m, m'\} = (mm'm^{-1})(\partial_1 m m')^{-1},$
2.  $\{\partial_2 l, \partial_2 l'\} = [l, l'] := ll'l^{-1}l'^{-1}$
3. (a)  $\{m, m'm''\} = \{m, m'\}^{mm'm^{-1}}\{m, m''\}$   
(b)  $\{mm', m''\} = \{m, m'm''m'^{-1}\}^{\partial_1 m}\{m', m''\},$
4.  $\{m, \partial_2 l\} = ({}^m l)(\partial_1 m l)^{-1}$
5.  ${}^n\{m, m'\} = \{{}^n m, {}^n m'\},$

where  $l, l' \in L$ ,  $m, m', m'' \in M$  and  $n \in N$ .

Here  ${}^m l$  denotes the action

$$\begin{aligned} M \times L &\rightarrow L \\ (m, l) &\mapsto {}^m l := l\{\partial_2 l^{-1}, m\}. \end{aligned} \tag{3}$$

A *normal* complex is one in which *im*  $\partial$  is normal in *ker* partial for all differentials.

It follows from these conditions that  $\partial_2 : L \rightarrow M$  is a crossed module with the action (3).

To get from a 3-group  $G_{(3)}$  to a 2-crossed module, we emulate the construction of a crossed module from a 2-group: one identifies

- $N$  is the group of objects of  $G_{(3)}$ .
- $M$  is the group of 1-morphisms of  $G_{(3)}$  starting at the identity object.
- $L$  is the group of 2-morphisms starting at the identity 1-arrow of the identity object
- $\partial_1 : M \rightarrow N$  is the target homomorphism such that  $m : \text{Id} \rightarrow \partial_1(m)$  for all  $m \in M$ .
- $\partial_2 : L \rightarrow M$  is the target homomorphism such that  $l : \text{Id}_{\text{Id}} \rightarrow \partial_2(l)$  for all  $l \in L$ .
- The various actions arise by whiskering, analogously to the case of a 2-group.

We will not go into the proof that this gives rise to a 2-crossed module for all 3-groups, but only in the case we are considering. One reason to consider 2-crossed modules is that the homotopy groups of  $G_{(3)}$  can be calculated as the homology of the sequence underlying the 2-crossed module.

### 2.3 Mapping cones of crossed modules

Another notion equivalent to 3-groups is crossed modules internal to crossed modules (more technically known as crossed squares, [10],[7]). More generally, consider a map  $\phi$  of crossed modules:

**Definition 4** (nonabelian mapping cone [7]). *For*

$$\begin{array}{ccc} H_2 & \xrightarrow{\phi_H} & H_1 \\ \downarrow t_2 & & \downarrow t_1 \\ G_2 & \xrightarrow{\phi_G} & G_1 \end{array}$$

a 2-term complex of crossed modules  $(t_i : H_i \rightarrow G_i)$ , we say its mapping cone is the complex of groups

$$H_2 \xrightarrow{\partial_2} G_2 \ltimes H_1 \xrightarrow{\partial_1} G_1, \quad (4)$$

where

$$\partial_1 : (g_2, h_1) \mapsto t_1(h_1)\phi_G(g_2)$$

and

$$\partial_2 : h_2 \mapsto (t_2(h_2), \phi_H(h_2)^{-1}).$$

Here  $G_2$  acts on  $H_1$  by way of the morphism  $\phi_G : G_2 \rightarrow G_1$ .

When no structure is imposed on  $\phi$ , (4) is merely a complex. However, if  $\phi$  is a crossed square, the mapping cone is a 2-crossed module (originally shown in [4], but see [5]). We will not need to define crossed squares here (the interested reader may consult [5]), but just note they come equipped with a map

$$h : G_2 \times H_1 \rightarrow H_2$$

satisfying conditions similar to the Peiffer lifting.

The only crossed square we will see in this paper is the identity map on a crossed module

$$\begin{array}{ccc} H & \xrightarrow{\text{id}} & H \\ \downarrow t & & \downarrow t \\ G & \xrightarrow{\text{id}} & G \end{array}$$

with the structure map

$$h : G \times H \rightarrow H \quad (5)$$

$$(g, h) \mapsto h^g h^{-1}. \quad (6)$$

This concept of “crossed modules of crossed modules” is explored in Norrie’s thesis [9] on ‘actors’ of crossed modules, with a focus on categorifying group theory, rather than geometry.

### 3 Main results

#### 3.1 The exact sequence $G_{(2)} \rightarrow \text{INN}_0(G_{(2)}) \rightarrow \Sigma G_{(2)}$

We describe the 3-group  $\text{INN}_0(G_{(2)})$  for  $G_{(2)}$  any strict 2-group, and show that it plays the role of the universal principal  $G_{(2)}$ -bundle in that

- $\text{INN}_0(G_{(2)})$  is equivalent to the trivial 3-group (hence “contractible”).
- $\text{INN}_0(G_{(2)})$  fits into the short exact sequence

$$G_{(2)} \hookrightarrow \text{INN}_0(G_{(2)}) \twoheadrightarrow \Sigma G_{(2)}$$

of strict 2-groupoids.

#### 3.2 $\text{INN}_0(G_{(2)})$ from a mapping cone

We show that the 3-group  $\text{INN}_0(G_{(2)})$  comes from a 2-crossed module

$$H \longrightarrow G \ltimes H \longrightarrow G$$

which is the mapping cone of

$$\begin{array}{ccc} H & \xrightarrow{\text{Id}} & H \\ t \downarrow & & \downarrow t \\ G & \xrightarrow{\text{Id}} & G \end{array},$$

the identity map of the crossed module  $(t : H \rightarrow G)$  which determines  $G_{(2)}$ .

Notice that this harmonizes with the analogous result for Lie 2-algebras discussed in [8].

### 4 Inner automorphism $n$ -groups

An automorphism of an  $n$ -group  $G_{(n)}$  is simply an automorphism of the  $n$ -category  $\Sigma G_{(n)}$ . We want to say that such an automorphism  $q$  is *inner* if it is

equivalent to the identity automorphism

$$\begin{array}{ccc}
 & \text{Id} & \\
 \nearrow & \Downarrow \sim & \searrow \\
 \Sigma G_{(n)} & & \Sigma G_{(n)} . \\
 \nwarrow & \Uparrow q & \nearrow
 \end{array}$$

A useful way to think of the  $n$ -groupoid of inner automorphisms is in terms of what we call “tangent categories”, a slight variation of the concept of comma categories.

Tangent categories in general happen to live in interesting exact sequences. In order to be able to talk about these, we first quickly set up our definitions for exact sequences of strict 2-groupoids.

Remember that we work entirely within the Gray category whose objects are strict 2-groupoids, whose morphisms are strict 2-functors, whose 2-morphisms are pseudonatural transformations and whose 3-morphisms are modifications of these.

#### 4.1 Exact sequences of strict 2-groupoids

Inner automorphism  $n$ -groups turn out to live in interesting *exact sequences of  $(n + 1)$ -groups*. Therefore we want to talk about generalizations of exact sequences of groups to the world of  $n$ -groupoids. Since for our purposes here only strict 2-groupoids matter, we shall be content with just using a definition applicable to that case.

**Definition 5** (exact sequence of strict 2-groupoids). *A collection of composable morphisms*

$$C_0 \xrightarrow{f_1} C_1 \xrightarrow{f_2} \cdots \xrightarrow{f_n} C_n$$

*of strict 2-categories  $C_i$  is called an exact sequence if, as ordinary maps between spaces of 2-morphisms,*

- $f_0$  is injective
- $f_n$  is surjective
- the image of  $f_i$  is the preimage under  $f_{i+1}$  of the collection of all identity 2-morphisms on identity 1-morphisms in  $\text{Mor}_2(C_{i+1})$ , for all  $1 \leq i < n$ .

In order to make this harmonize with our distinction between  $n$ -groups  $G_{(n)}$  and the corresponding 1-object  $n$ -groupoids  $\Sigma G_{(n)}$  we add to that

**Definition 6** (exact sequences of strict 2-groups). *A collection of composable morphisms*

$$G_0 \xrightarrow{f_1} G_1 \xrightarrow{f_2} \cdots \xrightarrow{f_n} G_n$$

of strict 2-groups is called an exact sequence if the corresponding chain

$$\Sigma G_0 \xrightarrow{\Sigma f_1} \Sigma G_1 \xrightarrow{\Sigma f_1} \cdots \xrightarrow{\Sigma f_n} \Sigma G_n$$

is an exact sequence of strict 2-groupoids.

**Remark.** Ordinary exact sequences of groups are thus precisely correspond to exact sequences of strict 2-groups all whose morphisms are identities.

## 4.2 Tangent 2-categories

We present a simple but useful way describe 2-categories of morphisms with coinciding source.

**Definition 7** (the point). *The point is the strict 2-category*

$$\mathbf{pt} := \{\bullet\}$$

with a single object and no nontrivial morphisms. We shall carefully distinguish this from the strict 2-groupoid

$$\mathbf{pt} := \{ \bullet \xrightarrow{\sim} \circ \},$$

consisting of two objects connected by a 1-isomorphism.

The 2-groupoid  $\mathbf{pt}$  might be called the “fat point”. It is of course equivalent to the point – but not isomorphic. We fix one injection

$$i : \mathbf{pt} \hookrightarrow \mathbf{pt}$$

$$i : \bullet \mapsto \bullet$$

once and for all.

It is useful to think of morphisms

$$\mathbf{f} : \mathbf{pt} \rightarrow C$$

from the fat point to some codomain  $C$  as labeled by the corresponding image of the ordinary point

$$\begin{array}{ccc} \mathbf{pt} & \xrightarrow{\mathbf{f}} & C \\ \downarrow & & \downarrow = \\ \mathbf{pt} & \xrightarrow{\mathbf{f}} & C \end{array}$$

**Definition 8** (tangent 2-bundle). *Given any strict 2-category  $C$ , we define its tangent 2-bundle*

$$TC \subset \text{Hom}_{2\text{Cat}}(\mathbf{pt}, C)$$

*to be that sub 2-category of morphisms from the fat point into  $C$  which collapses to a 0-category when pulled back along the fixed inclusion  $i : \mathbf{pt} \hookrightarrow \mathbf{pt} : \text{the morphisms } h \text{ in } TC \text{ are all those for which}$*

$$\begin{array}{ccc} \mathbf{pt} & & \mathbf{pt} \\ \downarrow & & \downarrow \\ \mathbf{pt} & \xrightarrow{\quad \mathbf{f} \quad} & C \\ & \Downarrow h & \\ & \mathbf{f}' & \end{array} = \begin{array}{ccc} \mathbf{pt} & & \mathbf{pt} \\ \downarrow & & \downarrow \\ \mathbf{pt} & \xrightarrow{\quad \mathbf{f} \quad} & C \end{array} .$$

*The tangent 2-bundle is a disjoint union*

$$TC = \bigoplus_{x \in \text{Obj}(C)} T_x C$$

*of tangent 2-categories at each object  $x$  of  $C$ . In this way it is a 2-bundle*

$$p : TC \longrightarrow \text{Obj}(C)$$

*over the space of objects of  $C$ .*

As befits a tangent bundle, the tangent 2-bundle has a canonical section

$$e_{\text{Id}} : \text{Obj}(C) \rightarrow TC$$

which sends every object of  $C$  to the Identity morphism on it.

**Example (slice categories).** For  $C$  any 1-groupoid, i.e. a strict 2-groupoid with only identity 2-morphisms, its tangent 1-category is the comma category


$$TC = ((\text{Obj}(C) \hookrightarrow C) \downarrow \text{Id}_C) .$$

This is the disjoint union of all co-over categories on all objects of  $C$

$$TC = \bigoplus_{a \in \text{Obj}(C)} (a \downarrow C)$$



Objects of  $TC$  are morphisms  $f : a \rightarrow b$  in  $C$ , and morphisms  $f \xrightarrow{h} f'$  in  $TC$  are commuting triangles




in  $C$ .


**Example (strict tangent 2-groupoids).** The example which we are mainly interested in is that where  $C$  is a strict 2-groupoid. For  $a$  any object in  $C$ , an object of  $T_a C$  is a morphism

$$a \xrightarrow{q} b.$$

A 1-morphism in  $T_a C$  is a filled triangle



in  $C$ . Finally, a 2-morphism in  $T_a C$  looks like



The composition of these 2-morphisms is the obvious one. We give a detailed description for the case the  $C = \Sigma G_{(2)}$  in 5.

**Proposition 1.** *For any strict 2-category  $C$ , its tangent 2-bundle  $TC$  fits into an exact sequence*

$$\text{Mor}(C) \hookrightarrow TC \twoheadrightarrow C$$

*of strict 2-categories.*

Here  $\text{Mor}(C) := \text{Disc}(\text{Mor}(C))$  is the 1-category of morphisms of  $C$ , regarded as a strict 2-category with only identity 2-morphisms.

Proof. The strict inclusion 2-functor on the left is

$$\left( g \xrightarrow{h} g' \right) \mapsto \begin{array}{ccc} & & b \\ & \nearrow g & \\ a & & \\ & \searrow g' & \\ & & b \end{array} \quad \begin{array}{c} \text{h} \\ \text{h} \\ \text{id} \end{array} \quad \begin{array}{c} \text{id} \\ \text{id} \\ \text{id} \end{array}$$

for  $g, g' : a \rightarrow b$  any two parallel morphisms in  $C$  and  $h$  any 2-morphism between them.

The strict surjection 2-functor on the right is

$$\begin{array}{ccc} & & b \\ & \nearrow q & \\ a & & \\ & \searrow k & \\ & & b' \end{array} \quad \begin{array}{c} F \\ K \\ f \end{array} \quad \begin{array}{c} L \\ L \\ k \end{array} \mapsto \begin{array}{ccc} & & b' \\ & \nearrow f & \\ b & & \\ & \searrow k & \\ & & b' \end{array} \quad \begin{array}{c} L \\ L \\ k \end{array}$$

The image of the injection is precisely the preimage under the surjection of the identity 2-morphism on the identity 1-morphisms. This means the sequence is exact.  $\square$

### 4.3 Inner automorphisms

Often, for  $G$  any group, inner and outer automorphisms are regarded as sitting in a short exact sequence

$$\text{Inn}(G) \longrightarrow \text{Aut}(G) \longrightarrow \text{Out}(G)$$

of ordinary groups.

But we will find shortly that we ought to be regarding the conjugation automorphisms by two group elements which differ by an element in the center of the group as *different* inner automorphisms.

So adopting this point of view for ordinary groups, one gets instead the exact sequence

$$\text{Z}(G) \longrightarrow \text{Inn}'(G) \longrightarrow \text{Aut}(G) \longrightarrow \text{Out}(G) .$$

Of course this means setting  $\text{Inn}'(G) \simeq G$ , which seems to make this step rather ill motivated. But it turns out that this degeneracy of concepts is a coincidence of low dimensions and will be lifted as we pass to inner automorphisms of higher groups.

First recall the standard definitions of center and automorphism of 2-groupoids:

**Definition 9.** *Given any strict 2-groupoid  $C$ ,*

- *the automorphism 3-group*

$$\text{AUT}(C) := \text{Aut}_{2\text{Cat}}(C)$$

*is the 2-groupoid of equivalences on  $C$ ;*

- *the center of  $C$*

$$Z(C) := \Sigma \text{AUT}(\text{Id}_C)$$

*is the (suspended) automorphism 2-group of the identity on  $C$ .*

**Example.** The automorphism 2-group of any ordinary group  $G$  (regarded as a 2-group  $\text{Disc}(G)$  with only identity morphisms)

$$\text{AUT}(G) := \text{AUT}(\Sigma G)$$

is that coming from the crossed module

$$G \xrightarrow{\text{Ad}} \text{Aut}(G) \xrightarrow{\text{Id}} \text{Aut}(G).$$

The center

$$Z(G) := Z(\Sigma G)$$

of any ordinary group is indeed the ordinary center of the group, regarded as a 1-object category.

To these two standard definitions, we add the following one, which is supposed to be the proper 2-categorical generalization of the concept of inner automorphisms.

**Definition 10** (inner automorphisms). *Given any strict 2-groupoid  $C$ , the tangent 2-groupoid*

$$\text{INN}(C) := T_{\text{Id}_C}(\text{Aut}_{2\text{Cat}}(C))$$

*is, called the 2-groupoid of inner automorphisms of  $C$ , and as such thought of as being equipped with the monoidal structure inherited from  $\text{End}(C)$ .*

If the transformation starting at the identity is denoted  $q$ , it makes good sense to call the inner automorphism being the target of that transformation  $\text{Ad}_q$ :

$$\begin{array}{ccc} & \text{Id} & \\ & \downarrow & \\ C & \begin{array}{c} \sim \\ \parallel \\ q \\ \downarrow \end{array} & C \\ & \text{Ad}_q & \end{array} .$$

A bigon of this form is an object in  $\text{INN}(C)$ . The product of two such objects is the horizontal composition of these bigons in  $2\text{Cat}$ . We shall spell this out in great detail for the case  $C = \Sigma G_{(2)}$  in 5.

**Proposition 2.** *For  $C$  any strict 2-category, we have canonical morphisms*

$$Z(C) \hookrightarrow \text{INN}(C) \longrightarrow \text{AUT}(C)$$

*of strict 2-categories whose composition sends everything to the identity 2-morphism on the identity 1-morphism on the identity automorphism of  $C$ .*

*Moreover, this sits inside the exact sequence from proposition 1 as*

$$\begin{array}{ccccc} Z(C) & \longrightarrow & \text{INN}(C) & \longrightarrow & \text{AUT}(C) \\ \downarrow & & \downarrow & & \downarrow \\ \text{Mor}(\mathbf{C}) & \longrightarrow & T\mathbf{C} & \longrightarrow & \mathbf{C} \end{array},$$

where  $\mathbf{C} := \text{Aut}_{2\text{Cat}}(C)$ .

Proof. Recall that a morphism in  $Z(C)$  is a transformation of the form

$$\begin{array}{ccc} & \text{Id} & \\ & \downarrow & \\ C & \xrightarrow{\sim} & C \\ & \downarrow q & \\ & \text{Ad}_q = \text{Id} & \end{array}.$$

This gives the obvious inclusion  $Z(G) \hookrightarrow \text{INN}(G)$ . The morphism  $\text{INN}(G) \rightarrow \text{AUT}(G)$  maps

$$\begin{array}{ccc} & \text{Id} & \\ & \downarrow & \\ C & \xrightarrow{\sim} & C \\ & \downarrow q & \\ & \text{Ad}_q & \end{array} \mapsto C \xrightarrow{\text{Ad}_q} C.$$

□

**Remark.** One would now want to define and construct the cokernel  $\text{OUT}(C)$  of the morphism  $\text{INN}(C) \rightarrow \text{AUT}(C)$  and then say that

$$Z(C) \longrightarrow \text{INN}(C) \longrightarrow \text{AUT}(C) \longrightarrow \text{OUT}(C)$$

is an exact sequence of 3-groups. But here we do not further consider this.

#### 4.4 Inner automorphism 3-groups.

Now we apply the general concept of inner automorphisms to 2-groups. The following definition just establishes the appropriate shorthand notation.

**Definition 11.** For  $G_{(2)}$  a strict 2-group, we write

$$\text{INN}(G_{(2)}) := \text{INN}(\Sigma G_{(2)})$$

for its 3-group of inner automorphisms.

In general this notation could be ambiguous, since one might want to consider the inner automorphisms of just the 1-groupoid underlying  $G_{(2)}$ . However, in the present context this will never occur and using the above definition makes a couple of expressions more manifestly appear as generalizations of familiar ones.

**Example.** For  $G$  an ordinary group, regarded as a discrete 2-group, one finds that

$$\text{INN}(G) := T_{\text{Id}_{\Sigma G}}(n\text{Cat}) \simeq T_{\bullet}(\Sigma G)$$

is the codiscrete groupoid over the elements of  $G$ . Its nature as a groupoid is manifest from its realization as

$$\text{INN}(G) = T_{\bullet}(\Sigma G).$$

But it is also a (strict) 2-group. The monoidal structure is that coming from its realization as  $\text{INN}(G) := T_{\text{Id}_{\Sigma G}}(n\text{Cat})$ . The crossed module corresponding to this strict 2-group is

$$G \xrightarrow{\text{Id}} G \xrightarrow{\text{Ad}} \text{Aut}(G).$$

The main point of interest for us is the generalization of this fact to strict 2-groups. One issue that one needs to be aware of then is that the above isomorphism  $T_{\text{Id}_{\Sigma G}}(n\text{Cat}) \simeq T_{\bullet}(\Sigma G)$  becomes a mere inclusion.

**Proposition 3.** For  $G_{(2)}$  any strict 2-group, we have an inclusion

$$T_{\bullet}\Sigma G_{(2)} \subset T_{\text{Id}_{\Sigma G_{(2)}}}(\text{Aut}_{2\text{Cat}}(\Sigma G_{(2)}))$$

of strict 2-groupoids.

This realizes  $T_{\bullet}\Sigma G_{(2)}$  as a sub 2-groupoid of  $T_{\text{Id}_{\Sigma G_{(2)}}}(\text{Aut}_{2\text{Cat}}(\Sigma G_{(2)}))$ .

*Proof.* The inclusion is essentially fixed by its action on objects: we define that an object in  $T_{\bullet}\Sigma G_{(2)}$ , which is a morphism

$$\bullet \xrightarrow{q} \bullet$$

in  $\Sigma G$ , is sent to the conjugation automorphism

$$\text{Ad}_q : \Sigma G_{(2)} \rightarrow \Sigma G_{(2)}$$

The transformation

$$\begin{array}{ccc} & \text{Id} & \\ \curvearrowright & \Downarrow q & \curvearrowleft \\ \Sigma G_{(2)} & & \Sigma G_{(2)} \\ & \text{Ad}_q & \end{array} .$$

connecting this to the identity is given by the component map

$$(\bullet \xrightarrow{g} \bullet) \mapsto \begin{array}{ccccc} \bullet & \xrightarrow{g} & \bullet & & \bullet \\ \downarrow q & & \downarrow q & \text{Id} & \downarrow q \\ \bullet & \xrightarrow{q^{-1}} \bullet & \xrightarrow{g} \bullet & \xrightarrow{q} \bullet & \bullet \end{array} .$$

In general one could consider transformations whose component maps involve here a non-identity 2-morphism. The inclusion we are describing picks out exactly those transformations whose component map only involves identity 2-morphisms.

The crucial point to realize now is the form of the component maps of morphisms

$$\begin{array}{ccc} & \text{Ad}_q & \\ \text{Id}_{\Sigma G_{(2)}} \curvearrowright & \Downarrow F & \downarrow \text{Ad}_F \\ & \text{Ad}_{q'} & \end{array} \quad \begin{array}{c} q \\ q' \end{array}$$

in  $T_{\text{Id}_{\Sigma G_{(2)}}}(\text{Aut}_{2\text{Cat}}(\Sigma G_{(2)}))$ .

The corresponding component map equation is

$$\begin{array}{ccc} \begin{array}{ccccc} \bullet & \xrightarrow{g} & \bullet & & \bullet \\ \downarrow q & & \downarrow q & \text{Id} & \downarrow q \\ \bullet & \xleftarrow{F} \bullet & \xrightarrow{\text{Ad}_q g} \bullet & & \bullet \\ \downarrow f & & \downarrow f & \text{Ad}_F(g) & \downarrow f \\ \bullet & \xrightarrow{\text{Ad}_{q'} g} \bullet & & & \bullet \end{array} & = & \begin{array}{ccccc} \bullet & \xrightarrow{g} & \bullet & & \bullet \\ \downarrow q' & & \downarrow q' & \text{Id} & \downarrow q' \\ \bullet & \xrightarrow{\text{Ad}_{q'} g} \bullet & & & \bullet \\ & & & & \downarrow f \\ & & & & \bullet \end{array} \end{array}$$

Solving this for  $\text{Ad}_F$  shows that this is given by conjugation

$$\text{Ad}_F : (\bullet \xrightarrow{g} \bullet) \mapsto \begin{array}{ccc} \bullet & \xrightarrow{q^{-1}} & \bullet \\ \downarrow f & \Downarrow & \downarrow f \\ \bullet & \xrightarrow{q'^{-1}} & \bullet \end{array} \xrightarrow{g} \begin{array}{ccc} \bullet & \xrightarrow{q} & \bullet \\ \downarrow f & \Downarrow F & \downarrow f \\ \bullet & \xrightarrow{q'} & \bullet \end{array}$$

with a morphism in  $T_\bullet(\Sigma G_{(2)})$ . And each such morphism in  $T_\bullet(\Sigma G_{(2)})$  yields a morphism in  $T_{\text{Id}_{\Sigma G_{(2)}}}(\text{Aut}_{2\text{Cat}}(\Sigma G_{(2)}))$  this way.

Finally, 2-morphisms in  $T_{\text{Id}_{\Sigma G_{(2)}}}(\text{Aut}_{2\text{Cat}}(\Sigma G_{(2)}))$  between these morphisms

$$\begin{array}{ccc} & \Sigma G_{(2)} & \\ \text{Ad}_q \swarrow & \text{Ad}_F & \searrow \text{Ad}_{q'} \\ & \Downarrow L & \\ & \Sigma G_{(2)} & \end{array}$$

come from component maps

$$\bullet \mapsto \begin{array}{ccc} & f & \\ & \Downarrow L & \\ & k & \end{array} \in \text{Mor}_2(\Sigma G_{(2)}) .$$

A sufficient condition for these component maps to solve the required condition for modifications of pseudonatural transformations is that they make

$$\begin{array}{ccc} & q & \\ & \Downarrow & \\ & F & \\ & K & \\ & f & \\ & \Downarrow L & \\ & k & \end{array}$$

2-commute. But this defines a 2-morphism in  $T_\bullet \Sigma G_{(2)}$ . And each such 2-morphism in  $T_\bullet(\Sigma G_{(2)})$  yields a 2-morphism in  $T_{\text{Id}_{\Sigma G_{(2)}}}(\text{Aut}_{2\text{Cat}}(C))$  this way.  $\square$

The crucial point is that by the embedding

$$T_\bullet \Sigma G_{(2)} \subset T_{\text{Id}_{\Sigma G_{(2)}}}(\text{Aut}_{2\text{Cat}}(\Sigma G_{(2)}))$$

the former 2-category inherits the monoidal structure of the latter and hence becomes a 3-group in its own right. This 3-group is the object of interest here.

## 5 The 3-group $\text{INN}_0(G_{(2)})$

**Definition 12** ( $\text{INN}_0(G_{(2)})$ ). *For  $G_{(2)}$  any strict 2-group, the 3-group  $\text{INN}_0(G_{(2)})$  is, as a 2-groupoid, given by*

$$\text{INN}_0(G_{(2)}) := T_\bullet \Sigma G_{(2)}$$

*and equipped with the monoidal structure inherited from the embedding of proposition 3.*

We now describe  $\text{INN}_0(G_{(2)})$  for  $G_{(2)}$  coming from the crossed module

$$H \xrightarrow{t} G \xrightarrow{\alpha} \text{Aut}(H)$$

in more detail, in particular spelling out the monoidal structure. We extract the operations in the crossed module corresponding to the various compositions in  $\text{INN}_0(G_{(2)})$  and then finally identify the 2-crossed module encoded by this.

### 5.1 Objects

The objects of  $\text{INN}_0(G_{(2)})$  are exactly the objects of  $G_{(2)}$ , hence the elements of  $G$ :

$$\text{Obj}(\text{INN}(G_{(2)})) = G.$$

The product of two objects in  $\text{INN}(G_{(2)})$  is just the product in  $G$ .

### 5.2 Morphisms

The morphisms

$$g \rightarrow h$$


in  $\text{INN}(G_{(2)})$  are

$$\begin{aligned} \text{Mor}(\text{INN}_0(G_{(2)})) &= \left\{ \begin{array}{c} \bullet \begin{array}{ccc} & \xrightarrow{g} & \bullet \\ & \parallel F & \downarrow f \\ & \xrightarrow{h} & \bullet \end{array} \end{array} \middle| \begin{array}{l} f, g, h \in G, F \in H \\ h = t(F)fg \end{array} \right\} \\ &= \{(f, F; g) \mid f, g \in G, F \in H\}. \end{aligned}$$




### 5.2.1 Composition

The composition of two such morphisms




is in terms of group labels given by




### 5.2.2 Product

Horizontal composition of automorphisms  $\Sigma G_{(2)} \rightarrow \Sigma G_{(2)}$  gives the product in the 3-group  $\text{INN}(G_{(2)})$

Left whiskering of pseudonatural transformations



amounts to the operation



on the corresponding triangles.

Right whiskering of pseudonatural transformations

$$\begin{array}{ccc}
 & \text{Ad}_q & \\
 \Sigma G_{(2)} & \begin{array}{c} \curvearrowright \\ \Downarrow \text{Ad}_F \\ \curvearrowleft \end{array} & \Sigma G_{(2)} \\
 & \text{Ad}_{q'} & \\
 & \xrightarrow{\text{Ad}_q} & \Sigma G_{(2)}
 \end{array}$$

amounts to the operation

$$\begin{array}{c} \bullet \end{array} \xrightarrow{q} \begin{array}{c} \bullet \\ \Downarrow F \\ \bullet \end{array} \xrightarrow{f} \begin{array}{c} \bullet \\ \Downarrow f \\ \bullet \end{array} \mapsto \begin{array}{c} \bullet \end{array} \xrightarrow{q} \begin{array}{c} \bullet \\ \Downarrow F \\ \bullet \end{array} \xrightarrow{f} \begin{array}{c} \bullet \\ \Downarrow f \\ \bullet \end{array} \xrightarrow{g} \begin{array}{c} \bullet \\ \Downarrow gfg^{-1} \\ \bullet \end{array} = \begin{array}{c} \bullet \end{array} \xrightarrow{gq} \begin{array}{c} \bullet \\ \Downarrow {}^g F \\ \bullet \end{array} \xrightarrow{gfg^{-1}} \begin{array}{c} \bullet \\ \Downarrow gfg^{-1} \\ \bullet \end{array}$$

on the corresponding triangles.

Since  $2\text{Cat}$  is a Gray category, the horizontal composition of pseudonatural transformations

$$\begin{array}{ccccc}
 & \text{Ad}_{q_1} & & \text{Ad}_{q_2} & \\
 \Sigma G_{(2)} & \begin{array}{c} \curvearrowright \\ \Downarrow \text{Ad}_{F_1} \\ \curvearrowleft \end{array} & \Sigma G_{(2)} & \begin{array}{c} \curvearrowright \\ \Downarrow \text{Ad}_{F_2} \\ \curvearrowleft \end{array} & \Sigma G_{(2)} \\
 & \text{Ad}_{q'_1} & & \text{Ad}_{q'_2} &
 \end{array}$$

is ambiguous. We shall agree to read this as

$$\begin{array}{ccccc}
 & \text{Ad}_{q_1} & & \text{Ad}_{q_2} & \\
 \Sigma G_{(2)} & \begin{array}{c} \curvearrowright \\ \Downarrow \text{Ad}_{F_1} \\ \curvearrowleft \end{array} & \Sigma G_{(2)} & \xrightarrow{\text{Ad}_{q_2}} & \Sigma G_{(2)} \\
 & \text{Ad}_{q'_1} & & \text{Ad}_{q'_2} & \\
 \Sigma G_{(2)} & \xrightarrow{\text{Ad}_{q'_1}} & \Sigma G_{(2)} & \begin{array}{c} \curvearrowright \\ \Downarrow \text{Ad}_{F_2} \\ \curvearrowleft \end{array} & \Sigma G_{(2)}
 \end{array} .$$


The corresponding operation on triangles labelled in the crossed module is

$$\begin{array}{c}
 \begin{array}{ccc}
 \bullet & \xrightarrow{q_1} & \bullet \\
 \Downarrow F_1 & & \downarrow f_1 \\
 \bullet & \xrightarrow{q_2} & \bullet \\
 \Downarrow F_2 & & \downarrow f_2 \\
 \bullet & & \bullet
 \end{array}
 & = &
 \begin{array}{ccc}
 \bullet & \xrightarrow{q_1} & \bullet \\
 \Downarrow F_1 & & \downarrow f_1 \\
 \bullet & \xrightarrow{q_2} & \bullet \\
 \Downarrow F_2 & & \downarrow f_2 \\
 \bullet & & \bullet
 \end{array}
 \\
 & &
 \begin{array}{ccc}
 \bullet & \xrightarrow{q_2 q_1} & \bullet \\
 \Downarrow q_2 F_1 & & \downarrow q_2 f_1 q_2^{-1} \\
 \bullet & \xrightarrow{\quad} & \bullet \\
 \Downarrow F_2 & & \downarrow f_2 \\
 \bullet & & \bullet
 \end{array}
 \\
 & &
 \begin{array}{ccc}
 \bullet & \xrightarrow{q_1} & \bullet \\
 \Downarrow F_2 f_2 q_2 F_1 & & \downarrow f_2 q_2 f_1 q_2^{-1} \\
 \bullet & & \bullet
 \end{array}
 \end{array}$$

The non-identitcal isomorphism which relates this to the other possible way to evaluate the horizontal composition of pseudonatural transformations gives rise to the Peiffer lifting of the corresponding 2-crossed module. This is discussed in 6.4.

### 5.3 2-Morphisms

The 2-morphisms in  $\text{INN}_0(G_{(2)})$  are given by diagrams




In terms of the group labels this means that  $L \in H$  satisfies


$$L = K^{-1}F. \quad (7)$$

#### 5.3.1 Composition

The horizontal composition of 2-morphisms in  $\text{INN}_0(G_{(2)})$  is given by




and vertical composition by




(Notice that these compositions do go horizontally and vertically, respectively, once we rotate such that the bigons have the standard orientation.)


Notice that whiskering along 1-morphisms




acts on the component maps as




and



There is one more type of whiskering possible with 2-morphisms,




which acts in the following way on the components:



where

$$\begin{aligned} F' &= F^f p G, \\ K' &= K^{kp} G, \\ L' &= {}^{kp} G^{-1} L^f p G. \end{aligned}$$


and



where


$$\begin{aligned} F' &= G^{gq}F, \\ K' &= G^{gq}K. \end{aligned}$$

An important case of this is:




### 5.3.2 Product

The whiskering along objects




gives the product of objects with 2-morphisms in the 3-group  $\text{INN}(G_{(2)})$ . Its action on 2-morphisms, which we have already discussed, extends in a simple way to 3-morphisms:


left whiskering along an object acts as



while right whiskering along an object acts as



To calculate the product of a pair of 2-morphisms, we use the fact that a 2-morphism is uniquely determined by its source and target.



with

$$\begin{aligned} F' &= F_2^{f_2 q_2} F_1 \\ K' &= K_2^{k_2 q_2} K_1 \\ L' &= L_2^{f_2 q_2} L_1 \end{aligned}$$

## 6 Properties of $\text{INN}_0(G_{(2)})$

### 6.1 Structure morphisms

We have defined  $\Sigma(\text{INN}_0(G_{(2)}))$  essentially as a sub 3-category of  $2\text{Cat}$ . The latter is a Gray category, in that it is a 3-category which is strict except for the exchange law for composition of 2-morphisms. Accordingly, also  $\Sigma(\text{INN}_0(G_{(2)}))$  is strict except for the exchange law for 2-morphisms.

This means that as a mere 2-groupoid (forgetting the monoidal structure)  $\text{INN}_0(G_{(2)})$  is strict.

#### 6.1.1 Strictness as a 2-groupoid

**Proposition 4.** *The underlying 2-groupoid of  $\text{INN}_0(G_{(2)})$  is strict.*



Proof. This follows from the rules for horizontal and vertical composition of 2-morphisms in  $\text{INN}_0(G_{(2)})$  – displayed in 5.3.1 – and the fact that  $G_{(2)}$  itself is a strict 2-group, by assumption.  $\square$


But the product 2-functor on  $\text{INN}_0(G_{(2)})$  respects horizontal composition in  $\text{INN}_0(G_{(2)})$  only weakly. In the language of 2-groups, this corresponds to a failure of the Peiffer identity

## 6.2 Trivializability

**Proposition 5.** *The 2-groupoid  $\text{INN}_0(G_{(2)})$  is connected,*

$$\pi_0(\text{INN}(G_1)) = 1.$$

Proof. For any two objects  $q$  and  $q'$  there is the morphism



$\square$

**Proposition 6.** *The Hom-groupoids of the 2-category  $\text{INN}_0(G_{(2)})$  are codiscrete, meaning that they have precisely one morphism for every ordered pair of objects.*

Proof. By equation (7) there is at most one 2-morphism between any parallel pair of morphisms in  $\text{INN}_0(G_{(2)})$ . For there to be any such 2-morphism at all, the two group elements  $f$  and  $k$  in the diagram above (7) have to satisfy  $kf^{-1} \in \text{im}(t)$ . But by using the source-target matching condition for  $F$  and  $K$  one readily sees that this is always the case.  $\square$

**Theorem 1.** *The 3-group  $\text{INN}_0(G_{(2)})$  is equivalent to the trivial 3-group. If  $G_{(2)}$  is a Lie 2-group, then  $\text{INN}_0(G_{(2)})$  is equivalent to the trivial Lie 3-group even as a Lie 3-group.*

Proof. Equivalence of 3-groups  $G_{(3)}, G'_{(3)}$  is, by definition, that of the corresponding 1-object 3-groupoids  $\Sigma G_{(3)}, \Sigma G'_{(3)}$ . For showing equivalence with the trivial 3-group, it suffices to exhibit a pseudonatural transformation of 3-functors

$$\text{id}_{\Sigma(\text{INN}_0(G_{(2)}))} \rightarrow I_{\Sigma(\text{INN}_0(G_{(2)}))},$$

where  $I_{\Sigma(\text{INN}_0(G_{(2)}))}$  sends everything to the identity on the single object of  $\Sigma \text{INN}_0(G_{(2)})$ . Such a transformation is obtained by sending the single object

to the identity 1-morphism on that object and sending any 1-morphism  $q$  to the 2-morphism  $q \rightarrow \text{id}$  from prop 5. By prop 6 this implies the existence of a unique assignment of a 3-morphism to any 2-morphism such that we do indeed obtain the component map of a pseudonatural transformation of 3-functors. By construction, this is clearly smooth when  $G_{(2)}$  is Lie.  $\square$

### 6.3 Universality

**Theorem 2.** *We have a short exact sequence of strict 2-groupoids*

$$G_{(2)} \hookrightarrow \text{INN}_0(G_{(2)}) \twoheadrightarrow \Sigma G_{(2)} .$$

Proof. This is just proposition 1, after noticing that

$$\text{Mor}(\Sigma G_{(2)}) = G_{(2)} .$$

$\square$

So the strict inclusion 2-functor on the left is

while the strict surjection 2-functor on the right is

### 6.4 The corresponding 2-crossed module

We now extract the structure of a 2-crossed module from  $\text{INN}_0(G_{(2)})$ . First, let

$$\text{Mor}_1^I = \text{Mor}_1(\text{INN}_0(G_{(2)}))|_{s^{-1}(\text{Id})}$$

and


$$\text{Mor}_2^I = \text{Mor}_2(\text{INN}_0(G_{(2)}))|_{s^{-1}(\text{id}_{\text{Id}})}$$

be subgroups of the 1- and 2-morphisms of  $\text{INN}_0(G_{(2)})$  respectively.


**Proposition 7.** *The group of 1-morphisms in  $\text{INN}_0(G_{(2)})$  starting at the identity object form the semidirect product group*

$$\text{Mor}_1^I = G \ltimes H$$

*under the identification*



*in that*




Proof. Use composition in  $\Sigma G_{(2)}$ . □

We have the obvious group homomorphism which is just the restriction of the target map

$$\partial_1 : \text{Mor}_1^I \rightarrow \text{Obj} := \text{Obj}(\text{INN}_0(G_{(2)}))$$

given by



This and the following constructions are to be compared with definition 4. There is an obvious action on  $\text{Mor}_1^I$ :

(8)

To see this, let

$$g = \partial_1 \left( \begin{array}{ccc} & \xrightarrow{\text{id}} & \bullet \\ & \parallel H & \downarrow h \\ \bullet & \searrow g & \bullet \end{array} \right) = t(H)h.$$

of the appropriate 2-cell considered as an element in the group  $\text{Mor}_1^I$  is

$$\left( \begin{array}{ccc} & \xrightarrow{\text{id}} & \bullet \\ \bullet & \begin{array}{c} \Downarrow H \\ \Downarrow \end{array} & \downarrow h \\ & \xrightarrow{t(H)h} & \bullet \end{array} \right)^{-1} = \begin{array}{ccc} & \xrightarrow{\text{id}} & \bullet \\ \bullet & \begin{array}{c} \Downarrow h^{-1} H^{-1} \\ \Downarrow \end{array} & \downarrow h^{-1} \\ & \xrightarrow{h^{-1} t(H)^{-1}} & \bullet \end{array} .$$


Therefore the conjugation is

$$=$$

(9)

10)

Though the Peiffer identity does not hold, both actions give rise to 2-cells with the same source and target, and hence define a 3-cell  $P$ . Denote the 2-cell (9) by  $(c, C; id)$  and the 2-cell (10) by  $(a, A; id)$  (for conjugation and action respectively).



Then


$$\begin{aligned}
P &= A^{-1}C \\
&= {}^gF^{-1} \left( H^h F^h f h^{-1} H^{-1} \right) \\
&= {}^gF^{-1} \left( {}^{t(H)}h F H^h f h^{-1} H^{-1} \right) \\
&= {}^gF^{-1} \left( {}^gF H^h f h^{-1} H^{-1} \right) \\
&= H^h f h^{-1} H^{-1}
\end{aligned}$$

However, what we really want is the Peiffer lifting, which will be a 3-cell with source the identity 2-cell. Hence,


**Proposition 8.** *The group of 2-morphisms in  $\text{INN}_0(G_{(2)})$  starting at the identity arrow on the identity object form the group*

$$\text{Mor}_2^I = H$$

*under the identification*




*in that*



Proof. Use the multiplication of 2-morphisms.  $\square$

So, we whisker the 3-cell  $(P; a, A; \text{id})$  above with the inverse of  $(a, A; \text{id})$ :



and the back face is necessarily  $P^{-1}$ .

**Definition 13** (Peiffer lifting). *Define the map*

$$\{\cdot, \cdot\} : \text{Mor}_1^I \times \text{Mor}_1^I \rightarrow \text{Mor}_2^I$$

by

$$\left\{ \begin{array}{c} \text{id} \\ \parallel \\ H \end{array} \right\} \left\{ \begin{array}{c} \text{id} \\ \parallel \\ F \end{array} \right\} = \begin{array}{c} \text{id} \\ \parallel \\ P \end{array} \quad t(P) \quad , \quad P = H^{hfh^{-1}}H^{-1}$$

Now define the homomorphism

$$\partial_2 : \text{Mor}_2^I \rightarrow \text{Mor}_1^I$$

by

which is again the restriction of the target map. Note there is an action of  $\text{Obj}$  on  $\text{Mor}_2^I$ :

$$\begin{array}{c}
 \bullet \xrightarrow{\text{id}} \bullet \xrightarrow{\text{id}} t(L) \xrightarrow{g} \bullet \xrightarrow{\text{id}} \bullet \xrightarrow{\text{id}} t(L) \xrightarrow{g} \bullet \xrightarrow{\text{id}} \bullet \xrightarrow{\text{id}} t(gL) \\
 \downarrow \text{id} \quad \downarrow \text{id} \quad \downarrow \text{id} \quad \downarrow \text{id} \quad \downarrow \text{id} \quad \downarrow \text{id} \quad \downarrow \text{id} \quad \downarrow \text{id} \\
 \bullet \xrightarrow{\text{id}} \bullet \xrightarrow{\text{id}} t(L) \xrightarrow{g} \bullet \xrightarrow{\text{id}} \bullet \xrightarrow{\text{id}} t(L) \xrightarrow{g} \bullet \xrightarrow{\text{id}} \bullet \xrightarrow{\text{id}} t(gL)
 \end{array}
 \quad (11)$$

Clearly  $\partial_2 \circ \partial_1$  is the constant map at the identity, and  $\text{im } \partial_2$  is a normal subgroup of  $\ker \partial_1$ , so

$$\mathrm{Mor}_2^I \xrightarrow{\partial_2} \mathrm{Mor}_1^I \xrightarrow{\partial_1} \mathrm{Obj} \quad (12)$$

is a sequence. We let the action of  $\text{Obj}$  on the other two groups be as described above in (8) and (11), and the maps  $\partial_2$  and  $\partial_1$  are clearly equivariant for this action.

**Proposition 9.** *The map  $\{\cdot, \cdot\}$  does indeed satisfy the properties of a Peiffer lifting, and (12) is a 2- crossed module.*

Proof. The first condition holds by definition, the second and the last one are easy to check. The others are tedious. It is easy, using the crossed module properties of  $H \rightarrow G$ , to calculate that the actions of  $\text{Mor}_1^I$  on  $\text{Mor}_2^I$  as defined from  $\text{INN}_0(G_{(2)})$  and as defined via  $\{\cdot, \cdot\}$  are the same.  $\square$

Since  $\text{im } \partial_2 = \ker \partial_1$ ,  $\partial_2$  is injective and  $\partial_1$  is onto, this shows that (12) has trivial homology and provides us with another proof that  $\text{INN}_0(G_{(2)})$  is contractible.



### 6.4.1 Relation to the mapping cone of $H \rightarrow G$

Given a crossed square

$$\begin{array}{ccc} L & \xrightarrow{f} & M \\ u \downarrow & & \downarrow v \\ N & \xrightarrow{g} & P \end{array}$$

with structure map  $h : N \times M \rightarrow L$ , Conduché [5] gives the Peiffer lifting of the mapping cone

$$L \longrightarrow N \ltimes M \longrightarrow P$$

as

$$\{(g, h), (k, l)\} = h(gkg^{-1}, h).$$

Recall from 2.3 that the identity map on  $t : H \rightarrow G$  is a crossed square with

$$h(g, h) = h^g h^{-1},$$

so the mapping cone is a 2-crossed module

$$H \xrightarrow{\partial_2} G \ltimes H \xrightarrow{\partial_1} G,$$

where

$$d_2(h) = (t(h), h^{-1}), \quad d_1(g, h) = t(h)g,$$

and with Peiffer lifting

$$\{(g_1, h_1), (g_2, h_2)\} = h_1^{g_1 g_2 g_1^{-1}} h_1^{-1}.$$

which is what we found for  $\text{INN}_0(G_{(2)})$ .

More precisely,

**Definition 14.** A morphism  $\psi$  of 2-crossed modules is a map of the underlying complexes

$$\begin{array}{ccccc} L_1 & \xrightarrow{\partial_2} & M_1 & \xrightarrow{\partial_1} & N_1 \\ \psi_L \downarrow & & \psi_M \downarrow & & \psi_N \downarrow \\ L_2 & \xrightarrow{\partial_2} & M_2 & \xrightarrow{\partial_1} & N_2 \end{array}$$

such that  $\psi_L$ ,  $\psi_M$  and  $\psi_N$  are equivariant for the  $N$ - and  $M$ -actions, and

$$\{\psi_M(\cdot), \psi_M(\cdot)\}_2 = \psi_L(\{\cdot, \cdot\}_1).$$

Using propositions 7 and 8, we have a map

$$\begin{array}{ccccc}
\mathrm{Mor}_2^I & \xrightarrow{\partial_2} & \mathrm{Mor}_1^I & \xrightarrow{\partial_1} & \mathrm{Obj} \\
\cong \downarrow & & \downarrow \cong & & \downarrow \cong \\
H & \xrightarrow{d_2} & G \ltimes H & \xrightarrow{d_1} & G
\end{array}$$

and the actions and Peiffer lifting agree, so

**Proposition 10.** *The 2-crossed module associated to  $\mathrm{INN}_0(G_{(2)})$  is isomorphic to the mapping cone of the identity map on the crossed module associated to  $G_{(2)}$ .*

---

## 7 Universal $n$ -bundles

In order to put the relevance of the 3-group  $\mathrm{INN}_0(G_{(2)})$  in perspective, we further illuminate our statement, 3.1, that  $\mathrm{INN}_0(G_{(2)})$  *plays the role of the universal  $G_{(2)}$ -bundle*. An exhaustive discussion will be given elsewhere.

### 7.1 Universal 1-bundles in terms of $\mathrm{INN}(G)$

Let  $\pi : Y \rightarrow X$  be a good cover of a space  $X$  and write  $Y^{[2]} := Y \times_X Y$  for the corresponding groupoid.

**Definition 15** ( $G$ -cocycles). *A  $G$ -(1-)cocycle on  $X$  is a functor*

$$g : Y^{[2]} \rightarrow \Sigma G.$$

This functor can be understood as arising from a choice

$$\pi^* P \xrightarrow[\sim]{t} Y \times G$$

of trivialization of a principal right  $G$ -bundle  $P \rightarrow X$  (which is essentially just a map to  $G$ ) as

$$g := \pi_2^* t \circ \pi_1^* t^{-1},$$

by noticing that  $G$ -equivariant isomorphisms

$$G \rightarrow G$$

are in bijection with elements of  $G$

$$g(x, y) : h \mapsto g(x, y)h$$

acting from the left.

**Observation 1** ( $G$ -bundles as morphisms of sequences of groupoids). *Given a  $G$ -cocycle on  $X$  as above, its pullback along the exact sequence*

$$G \longrightarrow \text{INN}(G) \longrightarrow \Sigma G ,$$

*which we write as*

$$\begin{array}{ccccccc} Y \times G & \longrightarrow & Y^{[2]} \times_g \text{INN}(G) & \longrightarrow & Y^{[2]} & \longrightarrow & X \\ \downarrow & & \downarrow & & \downarrow g & & \downarrow \\ G & \longrightarrow & \text{INN}(G) & \longrightarrow & \Sigma G & \longrightarrow & \{\bullet\} \end{array} ,$$

*produces the bundle of groupoids*

$$Y^{[2]} \times_g \text{INN}(G) \longrightarrow Y^{[2]}$$

*which plays the role of the total space of the  $G$ -bundle classified by  $g$ .*

This should be compared with the simplicial constructions described, for instance, in [6].

**Remark.** Using the fact that  $\text{INN}(G)$  is a 2-group, and using the injection  $G \rightarrow \text{INN}(G)$  we naturally obtain the  $G$ -action on  $Y^{[2]} \times_g \text{INN}(G)$ .

**Remark.** Notice that this is closely related to the integrated Atiyah sequence

$$\text{Ad}P \longrightarrow P \times_G P \longrightarrow X \times X$$

of groupoids over  $X \times X$  coming from the  $G$ -principal bundle  $P \rightarrow X$ :

$$\text{Ad}P \longrightarrow P \times_G P \longrightarrow X \times X$$

$$\begin{array}{ccccccc} Y \times G & \longrightarrow & Y^{[2]} \times_g \text{INN}(G) & \longrightarrow & Y^{[2]} & \longrightarrow & X \\ \downarrow & & \downarrow & & \downarrow g & & \downarrow \\ G & \longrightarrow & \text{INN}(G) & \longrightarrow & \Sigma G & \longrightarrow & \{\bullet\} \end{array} .$$

We now make precise in which sense, in turn,  $Y^{[2]} \times_g \text{INN}(G)$  plays the role of the total space of the  $G$ -bundle characterized by the cocycle  $g$ .

To reobtain the  $G$ -bundle  $P \rightarrow X$  from the groupoid  $Y^{[2]} \times_g \text{INN}(G)$  we form the pushout of

$$\begin{array}{ccc} Y^{[2]} \times_g \text{INN}(G) & \xrightarrow{\text{target}} & Y \times G \\ \text{source} \downarrow & & \\ & & Y \times G \end{array} \quad (13)$$

**Proposition 11.** *If  $g$  is the cocycle classifying a  $G$ -bundle  $P$  on  $X$ , then the pushout of 13 is (up to isomorphism) that very  $G$ -bundle  $P$ .*

Proof. Consider the square

$$\begin{array}{ccccc} Y^{[2]} \times_g \text{INN}(G) & \xrightarrow{\text{target}} & Y \times G & & \\ \text{source} \downarrow & & \downarrow t^{-1} & & \\ & & \pi^* P & \xrightarrow{t^{-1}} & P \\ & & \downarrow & & \\ Y \times G & \xrightarrow{t^{-1}} & \pi^* P & \longrightarrow & P \end{array} ,$$

where  $t : \pi^* P \xrightarrow{\sim} Y \times G$  is the local trivialization of  $P$  which gives rise to the transition function  $g$ . Then the diagram commutes by the very definition of  $g$ . Since  $t$  is an isomorphism and since  $\pi^* P \rightarrow P$  is locally an isomorphism, it follows that this is the universal pushout.  $\square$

## 7.2 Universal 2-bundles in terms of $\text{INN}_0(G_{(2)})$

Now let  $G_{(2)}$  be any strict 2-group. Let  $Y^{[3]}$  be the 2-groupoid whose 2-morphisms are triples of lifts to  $Y$  of points in  $X$ . A principal  $G_{(2)}$ -2-bundle [12, 13] has local trivializations characterized by 2-functors

$$g : Y^{[3]} \rightarrow \Sigma G_{(2)} .$$

**Definition 16** ( $G_{(2)}$ -cocycles). *A  $G_{(2)}$ -(2-)cocycle on  $X$  is a 2-functor*

$$g : Y^{[3]} \rightarrow \Sigma G_{(2)} .$$

(Instead of 2-functors on  $Y^{[3]}$  one could use pseudo functors on  $Y^{[2]}$ .)

As before, we can pull these back along our exact sequence of 2-groupoids

3.1

$$G_{(2)} \longrightarrow \text{INN}(G_{(2)}) \longrightarrow \Sigma G_{(2)}$$

to obtain

$$\begin{array}{ccccccc}
Y \times G_{(2)} & \longrightarrow & Y^{[3]} \times_g \text{INN}(G_{(2)}) & \longrightarrow & Y^{[3]} & \longrightarrow & X \\
\downarrow & & \downarrow & & \downarrow g & & \downarrow \\
G_{(2)} & \longrightarrow & \text{INN}(G_{(2)}) & \longrightarrow & \Sigma G_{(2)} & \longrightarrow & \{\bullet\}
\end{array} .$$

We reconstruct the total 2-space of the 2-bundle by forming the weak pushout of

$$\begin{array}{ccc}
Y^{[3]} \times_g \text{INN}(G_{(2)}) & \xrightarrow{\text{target}} & Y \times G_{(2)} \\
\text{source} \downarrow & & \\
Y \times G_{(2)} & & 
\end{array} . \tag{14}$$

Here “source” and “target” are defined relative to the inclusion

$$Y \times G_{(2)} \hookrightarrow Y^{[2]} \times_g \text{INN}(G_{(2)}) .$$


This means that for a given 1-morphism

$$\begin{array}{ccc}
\bullet & \xrightarrow{q_1} & \bullet \\
\Downarrow F & & \downarrow f := g(x, y) \\
\bullet & \xrightarrow{q_2} & \bullet
\end{array}
\quad
\begin{array}{c}
x \\
\downarrow \\
y
\end{array}$$


in  $Y^{[3]} \times_g \text{INN}(G_{(2)})$  (for any  $x, y \in Y$  with  $\pi(x) = \pi(y)$  and for  $g(x, y)$  the corresponding component of the given 2-cocycle) which we may equivalently rewrite as

$$\begin{array}{ccc}
\bullet & \xrightarrow{q_1} & \bullet \\
\Downarrow f^{-1} F & & \downarrow f \\
\bullet & \xrightarrow{f^{-1} q_2} & \bullet
\end{array}
\quad
\begin{array}{c}
x \\
\downarrow \\
y
\end{array}$$

the source in this sense is



regarded as a morphism in  $Y \times G_{(2)}$ , while the target is



regarded as a morphism in  $Y \times G_{(2)}$ .

This way the transition function  $g(x, y)$  acts on the copies of  $G_{(2)}$  which appear as the trivialized fibers of the  $G_{(2)}$ -bundle.

Bartels [12][proof of prop. 22] gives a reconstruction of total space of principal 2-bundle from their 2-cocycles which is closely related to  $Y^{[3]} \times_g \text{INN}(G_{(2)})$ .

We end by saying that  $\text{INN}(G_{(2)})$  is a special case of a much more general construction in 2-bundles [14], and that the results of this paper will, in a future paper, be connected to similar results using simplicial groups and simplicial universal bundles.

## References

- [1] J. Baez, A. Lauda, *Higher-Dimensional Algebra V: 2-Groups*, Th. and App. of Cat. 12 (2004), pp. 423–491, available as [math/0307200](#).
- [2] G. B. Segal, *Classifying spaces and spectral sequences*, Publ. Math. IHES No. 34 (1968) 105–112
- [3] P. Carrasco, A. M. Cegarra, *Group-theoretic algebraic models for homotopy types*, J. Pure Appl. Algebra **75** (1991) pp. 195–235.
- [4] D. Conduché, *Unpublished letter to R. Brown and J.-L. Loday*, 1984.
- [5] D. Conduché, *Simplicial crossed modules and mapping cones*, Georgian Mathematical Journal **10** (2003) pp. 623–636.
- [6] B. Jurčo, *Crossed Module Bundle Gerbes; Classification, String Group and Differential Geometry*, available as [math/0510078](#).

- [7] J.-L. Loday, *Spaces with finitely many homotopy groups*, J. Pure Appl. Algebra **24** (1982) pp. 179–202.
- [8] U. Schreiber, J. Stasheff, in preparation.
- [9] K. Norrie, *Actions and automorphisms of crossed modules*. Bulletin de la Société Mathématique de France, **118** no. 2 (1990), pp. 129–146.
- [10] D. Guin-Waléry and J.-L. Loday, *Obstructions à l’Excision en K-théorie Algébrique*, Springer Lecture Notes in Math., 854, (1981), pp. 179–216.
- [11] B. Day, R. Street, *Monoidal bicategories and Hopf algebroids*, Adv. Math **129** (1997) pp. 99–157.
- [12] T. Bartels, *2-Bundles*, PhD Thesis, available as [math/0410328](#).
- [13] I. Baković, *Bigroupoid bitorsors*, PhD Thesis.
- [14] D. M. Roberts, work in progress.
- [15] J. Baez, U. Schreiber, *Higher Gauge Theory*, in *Noncommutative Geometry and Representation Theory in Mathematical Physics*, Contemporary Mathematics 391, available as [math/0511710](#).
- [16] U. Schreiber, K. Waldorf, *Parallel transport and functors*, available as [arXiv:0705.0452](#).
- [17] J. Baez, D. Stevenson, *A classifying space for nonabelian cohomology*, in preparation