

COFINALITY

A CARDINAL κ IS SINGULAR
IFF

THE EXIST A CARDINAL $\lambda < \kappa$
AND SETS $\{S_\alpha : \alpha < \lambda\}$
SUCH THAT

$$- |S_\alpha| < \kappa \quad \text{ALL } \alpha$$

$$- \bigcup_{\alpha < \lambda} S_\alpha = \kappa.$$

THE LEAST SUCH λ IS $\text{CF } \kappa$.

[PROOF AND VARIATIONS
GROUP INTERACTION]

CARDINAL ARITHMETIC

HOW DOES γ DEPEND

ON α AND β

$$\text{IN } \aleph_\alpha^{\aleph_\beta} = \aleph_\gamma$$

WHICH γ GOES WITH
 α AND β ?

$$\aleph_0^{\aleph_0} = 2^{\aleph_0} = |\mathbb{R}| = \aleph_\gamma? \\ \equiv ?$$

$$\lceil \kappa + \lambda = | \kappa \times \{0\} \cup \lambda \times \{1\} |$$

$$\lfloor \kappa \cdot \lambda = | \underline{\kappa \times \lambda} |$$

$$\lfloor \text{FOR INFINITE } \kappa, \lambda : \max\{\kappa, \lambda\}$$

$$\Rightarrow \underset{\substack{\uparrow \\ \text{CARDINAL}}}{\kappa}^{\lambda} = | \underset{\substack{\uparrow \\ \text{SET OF FUNCTIONS}}}{\kappa}^{\lambda} | \quad (| {}^{\lambda}\kappa |)$$

NOTE: WE USE AC
TO GET $| {}^{\lambda}\kappa |$

IR AND SO ALSO $\mathcal{P}(\omega)$ NEED
NOT BE WELL-ORDERABLE.

$$2^{\aleph_0} = \dots \quad 2^{\aleph_1} = \dots$$

$$\bigcup_{2020}^{\aleph_{2021}} = \dots$$

$\rightarrow 2^{\aleph_0} = \aleph_1$ IS CANTOR'S
CONTINUUM HYPOTHESIS

$\rightarrow [\text{IF } X \in \mathbb{R} \text{ IS INFINITE}$
 $\text{THEN } |X| = |\mathbb{N}| \text{ OR } |X| = |\mathbb{R}|$
 CONSISTENT WITH & INDEP. OF ZFC

$$2^{\aleph_0} = \aleph_2, \dots, \aleph_{2020}, \dots$$

$$2^{\aleph_0} = \aleph_{\omega_1}, \dots$$

$$2^{\aleph_0} = \aleph_{\omega} \text{ IS } \underline{\underline{\text{FALSE}}}$$

WHAT CAN WE PROVE?

- λ INFINITE AND $2 \leq \kappa \leq 2^\lambda$
THEN $\kappa^\lambda = 2^\lambda$

$$2^\lambda \leq \kappa^\lambda : \text{EASY}$$

$$\kappa^\lambda \leq (2^\lambda)^\lambda \leq 2^{\lambda \cdot \lambda} = 2^\lambda$$

CANTOR - SCHRÖDER -
BERNSTEIN - DEDEKIND

$$S_0^{S_0} = S_1^{S_0} = \underline{\underline{(2^{S_0})^{S_0} = 2^{S_0}}}$$

APPARENTLY THINGS ARE
INTERESTING IF

$$\lambda < 2^\lambda < \kappa.$$

CERTAINLY:

$$\kappa \leq \kappa^\lambda \leq 2^\kappa$$

NON-TRIVIAL: $\kappa < \kappa^{\text{CF}\kappa}$

κ REG: $\kappa^\kappa = 2^\kappa$

LET $\langle \alpha_\eta : \eta < \text{CF}\kappa \rangle$ BE INCREASING
AND COFINAL IN κ .

LET $\langle f_\alpha : \alpha < \kappa \rangle$ BE
A SEQUENCE OF FUNCTIONS
FROM $\text{CF}\kappa$ TO κ .

IF $\eta < \text{cf } \kappa$ THEN

$$|\{f_\alpha(\eta) : \alpha < \alpha_\eta\}| \leq |\alpha_\eta| < \kappa.$$

DIAGONALISE DEFINE

$$f(\eta) = \min \kappa \setminus \{f_\alpha(\eta) : \alpha < \alpha_\eta\}$$

$f \neq f_\alpha$ FOR ALL α .

$$S_w^{S_0} > S_w$$

EVEN IF $2^{S_0} < S_w$

[SO IF $\text{cf } \kappa \leq \lambda < \kappa$
THEN $\kappa^\lambda > \kappa$.]

USEFUL: $[A]^\lambda = \{X \subseteq A : |X| = \lambda\}$

$$[\kappa]^\lambda = \kappa^\lambda \quad (\lambda \leq |A|)$$

$\geq$ κ^λ IS A SUBSET OF $[\lambda \times \kappa]^\lambda$
 $\uparrow$ FUNCTIONS $\lambda \leq \kappa \rightarrow \sim [\kappa]^\lambda$

$$\leq I = \{f \in \kappa^\lambda : f \text{ is 1-1}\}$$

$f \mapsto \text{RAN } f$ IS SURJECTIVE
FROM I TO $[\kappa]^\lambda$

AC SAYS: THERE IS AN
INJECTION $[\kappa]^\lambda \hookrightarrow I$.

$$|[\mathbb{R}]^{S_0}| = 2^{S_0}$$

SUMS/PRODUCTS OF SETS OF CARDINALS

$\langle \kappa_i : i < \lambda \rangle$ SEQUENCE OF CARDINALS.

ALL κ EQUAL: $\sum_{i < \lambda} \kappa_i = \kappa \cdot \lambda$


$$\sum_{i < \lambda} \kappa_i = \left| \bigcup_{i < \lambda} \{i\} \times \kappa_i \right|$$

IF λ IS INFINITE AND $\kappa_i > 0$ FOR ALL i THEN

$$\sum_{i < \lambda} \kappa_i = \lambda \cdot \sup_{i < \lambda} \kappa_i$$

$$\mu = \sup_{i < \lambda} \kappa_i$$

$$\leq: \bigcup_{i < \lambda} \{i\} \times \kappa_i \subseteq \lambda \times \mu$$



$$\geq: \lambda = \sum_{i < \lambda} 1 \leq \sum_{i < \lambda} \kappa_i$$

$$\kappa_j \leq \sum_{i < \lambda} \kappa_i \quad \text{FOR ALL } j$$

$$\mu \leq \sum_{i < \lambda} \kappa_i$$

$$\lambda \cdot \mu = \max\{\lambda, \mu\} \leq \sum_{i < \lambda} \kappa_i$$

$$\cdot \prod_{i < \lambda} \kappa_i = | \underbrace{\prod_{i < \lambda} \kappa_i}_{\text{CART. PRODUCT}} |$$

$\uparrow$
 CARDINAL

WHY CARDINAL AS INDEX SET

IF $f: J \rightarrow I$ IS A BIJECTION

$$\text{THEN } \prod_{i \in I} \kappa_i = \prod_{j \in J} \kappa_{f(j)}$$

$\uparrow$ $\uparrow$
 $\omega + 25$ ω

IF $\kappa_i \geq 2$ FOR ALL $i < \lambda$
 THEN

$$\sum_{i < \lambda} \kappa_i \leq \prod_{i < \lambda} \kappa_i$$

WHY $\kappa_i \geq 2$: $1 + 1 > 1 \cdot 1$
 ALSO THIS IS SHARP $2 + 2 = 2 \cdot 2$

λ FINITE : INDUCTION
 OTHERWISE

GIVEN $i < \lambda$ AND $\alpha \in \kappa_i$

PICK $f \in \prod_{i < \lambda} \kappa_i$ WITH $f(i) = \alpha$
 $\langle i, \alpha \rangle \mapsto \langle i, f \rangle$ INJECTIVE (CHECK)

FROM $\bigcup_{i < \lambda} \{i\} \times \kappa_i$ TO $\lambda \times \prod_{i < \lambda} \kappa_i$

$\downarrow$


$$\lambda < 2^\lambda \leq \prod_{i < \lambda} 2 \leq \prod_{i < \lambda} \kappa_i$$

SO $\lambda \cdot \prod_{i < \lambda} \kappa_i = \prod_{i < \lambda} \kappa_i$

PROJECT

DEFINE AN INJECTION
FROM

$$\bigcup_{i < \kappa} \{i\} \times \kappa_i \rightarrow \prod_{i < \kappa} \kappa_i$$



KÖNIG.

IF $\kappa_i < \lambda_i$ FOR $i < \mu$
THEN

$$\sum_{i < \mu} \kappa_i < \prod_{i < \mu} \lambda_i$$

WLOG $\kappa_i > 0$

SO $\lambda_i \geq 2$

$$\textcircled{1} \leq \sum_{i < \mu} \kappa_i \leq \sum_{i < \mu} \lambda_i \leq \prod_{i < \mu} \lambda_i$$

[EXPLICIT INJECTION POSSIBLE] $\lambda_i \geq 2$

$$\textcircled{2} \neq \text{LET } f: \bigcup_{i < \mu} \{i\} \times \kappa_i \rightarrow \prod_{i < \mu} \lambda_i$$

BE A MAP; WE SHOW f
IS NOT SURJECTIVE.

FIX j

$$\begin{aligned} \{j\} \times \kappa_j &\xrightarrow{f} \prod_{i < \mu} \lambda_i \xrightarrow{\pi_j} \lambda_j \\ (j, \alpha) &\mapsto f(j, \alpha) \mapsto f(j, \alpha)(j) \end{aligned}$$

THIS IS NOT ONTO

DEFINE $x \in \prod_{j < \mu} \lambda_j$

$$\text{BY } x(j) = \min \lambda_j \setminus \{f(j, \alpha)(j) : \alpha \in \kappa_j\}$$

THEN $x \notin \text{RAN } f$

CONSEQUENCES

$$- \kappa < 2^\kappa : \kappa_c = 1, \lambda_c = 2$$

$$- \kappa < \text{cf} 2^\kappa :$$

$$\text{TAKE } \kappa_c < 2^\kappa \text{ FOR } c < \kappa$$

$$\text{LET } \lambda_c = 2^\kappa$$

$$\sum_{c < \kappa} \kappa_c < \prod_{c < \kappa} \lambda_c = (2^\kappa)^\kappa = 2^\kappa$$

$$\text{CF } 2^{\aleph_0} > \aleph_1 \text{ THAT IS WHY } 2^{\aleph_0} \neq \aleph_\omega$$

$$- \kappa < \kappa^{\text{cf} \kappa}$$

$$\kappa = \sum_{c < \text{cf} \kappa} \kappa_c \quad \kappa_c < \kappa$$

$$\text{TAKE } \lambda_c = \kappa \quad c < \text{cf} \kappa$$

$$\kappa = \sum_{c < \text{cf} \kappa} \kappa_c$$

$$< \prod_{c < \text{cf} \kappa} \lambda_c = \kappa^{\text{cf} \kappa}$$

• THE CONTINUUM FUNCTION

$$\kappa \mapsto 2^\kappa$$

$$\aleph_\alpha \mapsto 2^{\aleph_\alpha} = \aleph_{\alpha+1}$$

GENERALIZED CONTINUUM HYPOTHESIS

$$2^{\aleph_\alpha} = \aleph_{\alpha+1} \quad \text{FOR ALL } \alpha.$$

• NOT FALSE [GÖDEL, 1940]

GCH IS CONSISTENT WITH ZFC

• NOT PROVABLE: [COHEN, 1963]

$$2^{\aleph_0} = \aleph_2 \text{ IS CONSISTENT.}$$

GCH MAKES LIFE EASY

- $2 \leq \kappa \leq 2^\lambda : \kappa^\lambda = 2^\lambda = \lambda^+$
- $\alpha \leq \beta : \sum_\alpha^{\beta+1} = \sum_{\beta+1}^\lambda$
- $\text{cf } \kappa \leq \lambda \leq \kappa : \kappa^\lambda = \kappa^+$

$$[\kappa < \kappa^\lambda \leq \kappa^\kappa = 2^\kappa = \kappa^+]$$
- $\lambda < \text{cf } \kappa : \kappa^\lambda = \kappa$

AS SETS $\kappa^\lambda = \bigcup_{\alpha < \kappa} \alpha^\lambda$

$$|\alpha^\lambda| = |\alpha|^\lambda \leq 2^{|\alpha| \cdot \lambda} = (|\alpha| \cdot \lambda)^+ \leq \kappa.$$

$$\kappa^\lambda \leq \sum_{\alpha < \kappa} |\alpha^\lambda| = \kappa \cdot \sup_{\alpha < \kappa} |\alpha|^\lambda = \kappa$$

BUT WHAT IN GENERAL?

- ⊗
- $\kappa < \lambda \rightarrow 2^\kappa \leq 2^\lambda$
 - $[2^{\aleph_0} = 2^{\aleph_1} \text{ is possible,}]$
 $\text{I.E., CONSISTENT WITH ZFC}]$
 - $\kappa < \text{cf } 2^\kappa$

⊗ THESE ARE THE ONLY RESTRICTIONS FOR REGULAR κ .

EASTON: IF $F: \text{CARD} \rightarrow \text{CARD}$ SATISFIES ⊗

THEN "FOR ALL REGULAR κ

$$F(\kappa) = 2^\kappa$$

IS CONSISTENT

κ LIMIT SO $\kappa = \sum_{i < \text{cf } \kappa} \kappa_i$

$$\boxed{2^\kappa = (2^{<\kappa})^{\text{cf } \kappa}}$$

$$2^{<\kappa} = \sup_{\lambda < \text{CARD}} \{2^\lambda : \lambda < \kappa\}$$

κ REGULAR: $2^\kappa = 2^\kappa$

$$2^\kappa = 2^{\sum \kappa_i} = \prod_i 2^{\kappa_i} \leq \prod_i 2^{<\kappa} = (2^{<\kappa})^{\text{cf } \kappa} \leq 2^\kappa$$

- IF κ IS SINGULAR AND THERE IS A $\mu < \kappa$ SUCH THAT

$$2^\nu = 2^\mu$$

FOR ALL ν WITH $\mu \leq \nu < \kappa$ THEN $2^\kappa = 2^\mu$.

WLOG $\mu \geq \text{cf} \kappa$.

$$- 2^{<\kappa} = 2^\mu$$

$$- 2^\kappa = (2^\mu)^{\text{cf} \kappa} \leq 2^{\mu \cdot \mu} = 2^\mu$$

GIMEL $\mathfrak{J}(\kappa) = \kappa^{\text{cf} \kappa}$.
(κ REGULAR $\mathfrak{J}(\kappa) = 2^\kappa$)

- IF FOR EVERY $\mu < \kappa$ THERE IS ν BETWEEN μ AND κ WITH $2^\nu > 2^\mu$ THEN $\text{cf}(2^{<\kappa}) = \text{cf} \kappa$.
THEN

$$\begin{aligned} 2^\kappa &= (2^{<\kappa})^{\text{cf} \kappa} \\ &= \mathfrak{J}(2^{<\kappa}) \end{aligned}$$

SUMMARY FOR 2^κ :

- κ SUCCESSOR: $2^\kappa = \kappa^\kappa = \mathfrak{J}(\kappa)$
- κ LIMIT 2^ν CONSTANT ON A TAIL BELOW κ :
 $2^\kappa = 2^{<\kappa} \cdot \mathfrak{J}(\kappa)$.
 - κ REGULAR: $\mathfrak{J}(\kappa) = 2^\kappa$
 - κ SINGULAR $2^{<\kappa} = 2^\kappa = \kappa^\kappa \geq \mathfrak{J}(\kappa)$
- κ LIMIT 2^ν NOT CONSTANT ON A TAIL:

$$2^\kappa = \mathfrak{J}(2^{<\kappa})$$

GIMEL IS IMPORTANT.

$$\aleph_\omega \text{ STRONG LIMIT} \rightarrow 2^{\aleph_\omega} = \aleph_{\omega_1}^{\aleph_\omega} < \aleph_{\omega_2}$$

$$2^{<\kappa} = \kappa$$

BACK TO EXPONENTIATION.

$$\kappa^{<\lambda} = \sup \{ \kappa^\mu : \mu < \lambda, \mu \text{ CARD} \}$$

(IF $\lambda = \mu^+$ THIS IS JUST κ^μ)

USEFUL FACT IF λ IS AN INFINITE CARDINAL AND $\langle \kappa_i : i < \lambda \rangle$ IS NON-DECREASING ($i < j \rightarrow \kappa_i \leq \kappa_j$)

$$\text{THEN } \prod_{i < \lambda} \kappa_i = (\sup_{i < \lambda} \kappa_i)^\lambda$$

[HOMEWORK]

TOWARDS A SUMMARY FOR κ^λ .

- IF κ IS REGULAR AND $\lambda < \kappa$ THEN $\kappa^\lambda = \bigcup_{\alpha < \kappa} \alpha^\lambda$ (SETS) AND SO

$$\kappa^\lambda = \sum_{\alpha < \kappa} |\alpha|^\lambda \quad (\text{CARDS})$$

- HAUSDORFF: $\aleph_{\alpha+1}^{\aleph_\beta} = \underbrace{\aleph_\alpha^{\aleph_\beta}}_{\text{MAXIMUM TERM}} \cdot \underbrace{\aleph_{\alpha+1}}_{\text{NUMBER OF TERMS}}$

$$\beta \leq \alpha$$

$$\beta > \alpha \quad \aleph_{\alpha+1}^{\aleph_\beta} \ll 2^{\aleph_\beta}$$

$$\lim_{\alpha \rightarrow \kappa} \alpha^\lambda = \sup \{ \mu^\lambda : \mu < \kappa, \mu \text{ CARD} \}$$

IF κ IS A LIMIT AND $\lambda \geq \text{cf} \kappa$
 THEN

$$\kappa^\lambda = \left(\lim_{\alpha \rightarrow \kappa} \alpha^\lambda \right)^{\text{cf} \kappa}$$

$$\kappa = \sum_{i < \text{cf} \kappa} \kappa_i \quad \text{with } 1 \leq \kappa_i < \kappa$$

$$\begin{aligned} \kappa^\lambda &\leq \left(\prod_{i < \text{cf} \kappa} \kappa_i \right)^\lambda = \prod_{i < \text{cf} \kappa} \kappa_i^\lambda \\ &\leq \prod_{i < \text{cf} \kappa} \left(\lim_{\alpha \rightarrow \kappa} \alpha^\lambda \right) \\ &= \left(\lim_{\alpha \rightarrow \kappa} \alpha^\lambda \right)^{\text{cf} \kappa} \\ &\leq (\kappa^\lambda)^{\text{cf} \kappa} \\ &= \kappa^\lambda \end{aligned}$$

SUMMARY: κ^λ

- $2 \leq \kappa \leq 2^\lambda$: $\kappa^\lambda = 2^\lambda$
- IF THERE IS $\mu < \kappa$ WITH $\kappa \leq \mu^\lambda$
 THEN $\kappa^\lambda = \mu^\lambda$
- IF $\kappa > \lambda$ AND $\mu^\lambda < \kappa$ FOR ALL $\mu < \kappa$
 THEN

$$\lambda \geq \text{cf} \kappa : \kappa^\lambda = \sqrt[\text{cf} \kappa]{\kappa^{\text{cf} \kappa}} \leftarrow \text{BECAUSE } \lim_{\alpha \rightarrow \kappa} \alpha^\lambda = \kappa$$

$$\lambda < \text{cf} \kappa : \kappa^\lambda = \kappa$$

$$\kappa = \mu^+ : \text{HANS DORFF}$$

$$\kappa^\lambda = \mu^\lambda \cdot \kappa$$

$$\kappa \text{ LIMIT } \kappa^\lambda = \lim_{\alpha \rightarrow \kappa} \alpha^\lambda = \kappa$$

$$\text{SETS: } \kappa^\lambda = \bigcup_{\alpha < \kappa} \alpha^\lambda$$

$$\bullet \kappa^\lambda \text{ CAN BE } \dots 2^\lambda$$

$$\dots \kappa$$

$$\dots \beth(M) \text{ FOR SOME } M \text{ WITH } \text{cf} M \leq \lambda < M.$$

THAT $\mu = \min\{\nu < \kappa : \nu^{\lambda} \geq \kappa\}$
 (IN THE CASE THAT $2^{\lambda} < \kappa$)

SINGULAR CARDINALS Hypothesis
 FOR EVERY SINGULAR κ

IF $2^{\text{cf} \kappa} < \kappa$ THEN $\kappa^{\text{cf} \kappa} = \kappa^{+}$

SCH SAYS

IF $2^{S_0^{+}} < S_{\omega}^{+}$

THEN $S_{\omega}^{+} = S_{\omega+1}^{+}$

[illegible]

[illegible]