

Some properties of the Thom spectrum over loop suspension of complex projective space

Andrew Baker and Birgit Richter

ABSTRACT. This note provides a reference for some properties of the Thom spectrum $M\xi$ over $\Omega\Sigma\mathbb{C}P^\infty$. Some of this material is used in recent work of Kitchloo and Morava. We determine the $M\xi$ -cohomology of $\mathbb{C}P^\infty$ and show that $M\xi^*(\mathbb{C}P^\infty)$ injects into power series over the algebra of non-symmetric functions. We show that $M\xi$ gives rise to a commutative formal group law over the non-commutative ring $\pi_*M\xi$. We also discuss how $M\xi$ and some real and quaternionic analogues behave with respect to spectra that are related to these Thom spectra by splittings and by maps.

Introduction

The map $\mathbb{C}P^\infty = BU(1) \rightarrow BU$ gives rise to a canonical loop map $\Omega\Sigma\mathbb{C}P^\infty \rightarrow BU$. Therefore the associated Thom spectrum has a strictly associative multiplication. But as is visible from its homology, which is a tensor algebra on the reduced homology of $\mathbb{C}P^\infty$, it is not even homotopy commutative. This homology ring coincides with the ring of non-symmetric functions, \mathbf{NSymm} . We show that there is a map from the $M\xi$ -cohomology of $\mathbb{C}P^\infty$ to the power series over the ring of non-symmetric functions, \mathbf{NSymm} . This result is used in [MK] in an application of $M\xi$ to quasitoric manifolds.

Although $M\xi$ maps to MU , there is no obvious map to it from MU , so a priori it is unclear whether there is a formal group law associated to $M\xi$. However, analogues of the classical Atiyah-Hirzebruch spectral sequence calculations for MU can be made for $M\xi$, and these show that there is a ‘commutative formal group’ structure related to $M\xi^*\mathbb{C}P^\infty$, even though the coefficient ring $M\xi_*$ is not commutative and its elements do not commute with the variables coming from the choices of complex orientations. A formal group law in this context is an element $F(x, y) \in M\xi^*(\mathbb{C}P^\infty \times \mathbb{C}P^\infty)$ that satisfies the usual axioms for a commutative formal group law. We describe the precise algebraic structure arising here in Section 3.

For MU the p -local splitting gives rise to a map of ring spectra $BP \rightarrow MU_{(p)}$. We show that despite the fact that $M\xi_{(p)}$ splits into (suspended) copies of BP , there is no map of ring spectra $BP \rightarrow M\xi_{(p)}$. For the canonical Thom spectrum over $\Omega\Sigma\mathbb{R}P^\infty$, $M\xi_{\mathbb{R}}$, we show that the map of E_2 -algebra spectra $HF_2 \rightarrow MO$ does not give rise to a ring map $HF_2 \rightarrow M\xi_{\mathbb{R}}$.

2010 *Mathematics Subject Classification.* Primary 55P43, Secondary 55N22.

For a map of ring spectra $MU \rightarrow E$ to some commutative S -algebra E one can ask whether a map of commutative S -algebras $S \wedge_{\mathbb{P}(S)} \mathbb{P}(\Sigma^{\infty-2}\mathbb{C}P^\infty) \rightarrow E$ factors over MU . Here $\mathbb{P}(-)$ denotes the free commutative S -algebra functor. It is easy to see that MU is not equivalent to $S \wedge_{\mathbb{P}(S)} \mathbb{P}(\Sigma^{\infty-2}\mathbb{C}P^\infty)$ and we show that there are commutative S -algebras for which such a factorization does not exist. In the associative setting, the analogous universal gadget would be $S \amalg_{\mathbb{A}(S)} \mathbb{A}(\Sigma^{\infty-2}\mathbb{C}P^\infty)$, where $\mathbb{A}(-)$ is the free associative S -algebra functor and \amalg denotes the coproduct in the category of associative S -algebras. It is obvious that the homology of $S \amalg_{\mathbb{A}(S)} \mathbb{A}(\Sigma^{\infty-2}\mathbb{C}P^\infty)$ is much bigger than the one of $M\xi$. If we replace the coproduct by the smash product, there is still a canonical map $S \wedge_{\mathbb{A}(S)} \mathbb{A}(\Sigma^{\infty-2}\mathbb{C}P^\infty) \rightarrow M\xi$ due to the coequalizer property of the smash product. However, this smash product still has homology that is larger than that of $M\xi$. Therefore the freeness of $\Omega\Sigma\mathbb{C}P^\infty$ is not reflected on the level of Thom spectra.

1. The Thom spectrum of ξ

Lewis showed [LMSM, Theorem IX.7.1] that an n -fold loop map to BF gives rise to an E_n -structure on the associated Thom spectrum. Here E_n is the product of the little n -cubes operad with the linear isometries operad. For a more recent account in the setting of symmetric spectra see work of Christian Schlichtkrull [Sch].

The map $j: \Omega\Sigma\mathbb{C}P^\infty \rightarrow BU$ is a loop map and so the Thom spectrum $M\xi$ associated to that map is an A_∞ ring spectrum and the natural map $M\xi \rightarrow MU$ is one of A_∞ ring spectra, or equivalently of S -algebras in the sense of [EKMM]. Since the homology $H_*(M\xi)$ is isomorphic as a ring to $H_*(\Omega\Sigma\mathbb{C}P^\infty)$, we see that $M\xi$ is not even homotopy commutative. We investigated some of the properties of $M\xi$ in [BR].

For any commutative ring R , under the Thom isomorphism $H_*(\Omega\Sigma\mathbb{C}P^\infty; R) \cong H_*(M\xi; R)$, the generator Z_i corresponds to an element $z_i \in H_{2i}(M\xi; R)$, where we set $z_0 = 1$. Thomifying the map $i: \mathbb{C}P^\infty \rightarrow \Omega\Sigma\mathbb{C}P^\infty$, we obtain a map $Mi: MU(1) \rightarrow \Sigma^2 M\xi$, and it is easy to see that

$$(1.1) \quad Mi_*\beta_{i+1} = z_i.$$

1.1. Classifying negatives of bundles. For every based space X , time-reversal of loops is a loop-map from $(\Omega X)^{\text{op}}$ to ΩX , *i.e.*,

$$(\bar{\cdot}): (\Omega X)^{\text{op}} \rightarrow \Omega X; \quad w \mapsto \bar{w},$$

where $\bar{w}(t) = w(1-t)$. Here $(\Omega X)^{\text{op}}$ is the space of loops on X with the opposite multiplication of loops.

We consider BU with the H -space structure coming from the Whitney sum of vector bundles and denote this space by BU_\oplus . A complex vector bundle of finite rank on a reasonable space Y is represented by a map $f: Y \rightarrow BU$ and the composition

$$Y \xrightarrow{f} BU_\oplus^{\text{op}} \xrightarrow{(\bar{\cdot})} BU_\oplus$$

classifies the negative of that bundle, switching the rôle of stable normal bundles and stable tangent bundles for smooth manifolds.

For line bundles $g_i: Y \rightarrow \mathbb{C}P^\infty \rightarrow BU_\oplus$ ($i = 1, \dots, n$) we obtain a map

$$g = (g_n, \dots, g_1): Y \rightarrow Y^n \rightarrow (\mathbb{C}P^\infty)^n \rightarrow (\Omega\Sigma\mathbb{C}P^\infty)^{\text{op}} \rightarrow BU_\oplus^{\text{op}}$$

and the composition with loop reversal classifies the negative of the sum $g_n \oplus \dots \oplus g_1$ as indicated in the following diagram.

$$\begin{array}{ccccc}
 & & g & & \\
 & \curvearrowright & & \curvearrowleft & \\
 Y & \xrightarrow{\quad} & Y^n & \longrightarrow & (\Omega\Sigma\mathbb{C}P^\infty)^{\text{op}} & \longrightarrow & BU_\oplus^{\text{op}} \\
 & \searrow & & & \downarrow (\bar{\cdot}) & & \downarrow (\bar{\cdot}) \\
 & & & & \Omega\Sigma\mathbb{C}P^\infty & \longrightarrow & BU_\oplus \\
 & \curvearrowleft & & \curvearrowright & & & \\
 & & g & & & &
 \end{array}$$

In this way, the splitting of the stable tangent bundle of a toric manifold into a sum of line bundles can be classified by $\Omega\Sigma\mathbb{C}P^\infty$. For work on an interpretation of $\pi_*M\xi$ as the habitat for cobordism classes of quasitoric manifolds see [MK].

2. $M\xi$ -(co)homology

We note that the composition of the natural map $i: \mathbb{C}P^\infty \rightarrow \Omega\Sigma\mathbb{C}P^\infty$ with $j: \Omega\Sigma\mathbb{C}P^\infty \rightarrow BU$ classifies the reduced line bundle $\eta-1$ over $\mathbb{C}P^\infty$. The associated map $Mi: \Sigma^\infty MU(1) \rightarrow \Sigma^2 M\xi$ gives a distinguished choice of complex orientation $x_\xi \in \widetilde{M\xi}^2(\mathbb{C}P^\infty)$, since the zero-section $\mathbb{C}P^\infty \rightarrow MU(1)$ is an equivalence.

We use the Atiyah-Hirzebruch spectral sequence

$$(2.1) \quad E_2^{*,*} = H^*(\mathbb{C}P^\infty; M\xi^*) \implies M\xi^*(\mathbb{C}P^\infty).$$

As $M\xi$ is an associative ring spectrum, this spectral sequence is multiplicative and its E_2 -page is $\mathbb{Z}[[x]] \otimes M\xi^*$. As the spectral sequence collapses, the associated graded is of the same form and we can deduce the following:

LEMMA 2.1. *As a left $M\xi^* = M\xi_{-*}$ -module we have*

$$(2.2) \quad M\xi^*(\mathbb{C}P^\infty) = \left\{ \sum_{i \geq 0} a_i x_\xi^i : a_i \in M\xi_*^* \right\}.$$

The filtration in the spectral sequence (2.1) comes from the skeleton filtration of $\mathbb{C}P^\infty$ and corresponds to powers of the augmentation ideal $\widetilde{M\xi}^*(\mathbb{C}P^\infty)$ in $M\xi^*(\mathbb{C}P^\infty)$. Of course the product structure in the ring $M\xi^*(\mathbb{C}P^\infty)$ is more complicated than in the case of $MU^*(\mathbb{C}P^\infty)$ since x_ξ is not a central element.

In order to understand a difference of the form $ux_\xi^k - x_\xi^k u$ with $u \in M\xi_m$ and $k \geq 1$ we consider the cofibre sequence

$$\Sigma^m \mathbb{C}P^{k-1} \subseteq \Sigma^m \mathbb{C}P^k \rightarrow \Sigma^m \mathbb{S}^{2k}.$$

Both elements ux_ξ^k and $x_\xi^k u$ restrict to the trivial map on $\Sigma^m \mathbb{C}P^{k-1}$. The orientation x_ξ restricted to \mathbb{S}^2 is the 2-fold suspension of the unit of $M\xi$, $\Sigma^2 i \in M\xi^2(\mathbb{S}^2)$.

Centrality of the unit ensures that the following square and outer diagram

$$\begin{array}{ccc}
 \Sigma^m S \wedge (\Sigma^2 S)^{\wedge k} & \xrightarrow{u \wedge (\Sigma^2 i)^{\wedge k}} & M\xi \wedge (\Sigma^2 M\xi)^{\wedge k} \\
 \downarrow \text{twist} & & \downarrow \text{twist} \\
 (\Sigma^2 S)^{\wedge k} \wedge \Sigma^m S & \xrightarrow{(\Sigma^2 i)^{\wedge k} \wedge u} & (\Sigma^2 M\xi)^{\wedge k} \wedge M\xi
 \end{array}
 \begin{array}{c}
 \\
 \nearrow \\
 \searrow \\
 \Sigma^{2k} M\xi
 \end{array}$$

commute, so the difference $ux_\xi^k - x_\xi^k u$ is trivial. This yields

LEMMA 2.2. *For every $u \in M\xi_m$ and $k \geq 1$, u and x_ξ^k commute up to elements of filtration at least $2k + 2$, i.e.,*

$$(2.3) \quad ux_\xi^k - x_\xi^k u \in (\widetilde{M\xi}^*(\mathbb{C}P^\infty))^{[2k+2]}.$$

Let E be any associative S -algebra with an orientation class $u_E \in E^2(\mathbb{C}P^\infty)$. The Atiyah-Hirzebruch spectral sequence for $E^*\mathbb{C}P^\infty$ identifies $E^*(\mathbb{C}P^\infty)$ with the left E_* -module of power series in u_E as in the case of $M\xi$:

$$E^*(\mathbb{C}P^\infty) = \left\{ \sum_{i \geq 0} \theta_i u_E^i : \theta_i \in E_* \right\}.$$

The orientation class $u_E \in E^2(\mathbb{C}P^\infty)$ restricts to the double suspension of the unit of E , $\Sigma^2 i_E \in E^2(\mathbb{C}P^1)$. Induction on the skeleta shows that for all n , $E_*(\mathbb{C}P^n)$ is free over E^* and we obtain that

$$E_*(\mathbb{C}P^\infty) \cong E_*\{\beta_0, \beta_1, \dots\}$$

with $\beta_i \in E_{2i}(\mathbb{C}P^\infty)$ being dual to u_E^i . Let $\varphi: M\xi \rightarrow H \wedge M\xi$ be the map induced by the unit of $H = H\mathbb{Z}$, and let $\Theta: M\xi^*(\mathbb{C}P^\infty) \rightarrow (H \wedge M\xi)^*(\mathbb{C}P^\infty)$ be the induced ring homomorphism (in fact Θ is a monomorphism as explained below). Then

$$(2.4) \quad \Theta(x_\xi) = \sum_{i \geq 0} z_i x_H^{i+1} = z(x_H),$$

where $x_H \in (H \wedge M\xi)^2(\mathbb{C}P^\infty)$ is the orientation coming from the canonical generator of $H^2(\mathbb{C}P^\infty)$. The proof is analogous to that for MU in [Ad]. Note that $H \wedge M\xi$ is an algebra spectrum over the commutative S -algebra H which acts centrally on $H \wedge M\xi$. Hence x_H is a central element of $(H \wedge M\xi)^*(\mathbb{C}P^\infty)$. This contrasts with the image of x_ξ in $(H \wedge M\xi)^*(\mathbb{C}P^\infty)$ which does not commute with all elements of $(H \wedge M\xi)_*$.

We remark that the cohomology ring $M\xi^*(\mathbb{C}P^\infty)$ is highly non-commutative. Using (2.4), and noting that coefficient $z_i \in H_{2i}(M\xi)$ is an indecomposable in the algebra $H_*(M\xi)$, it follows that x_ξ does not commute with any of the z_i . For example, the first non-trivial term in the commutator

$$z_1 z(x_H) - z(x_H) z_1$$

is $(z_1 z_2 - z_2 z_1) x_H^3 \neq 0$.

Let NSymm denote the ring of non-symmetric functions. This ring can be identified with $H_*(\Omega\Sigma\mathbb{C}P^\infty)$. Using this and the above orientation we obtain

PROPOSITION 2.3. *The map Θ induces a monomorphism*

$$\Theta: M\xi^*(\mathbb{C}P^\infty) \rightarrow \text{NSymm}[[x_H]].$$

PROOF. The right-hand side is isomorphic to the $H \wedge M\xi$ -cohomology of $\mathbb{C}P^\infty$. As $M\xi$ is a wedge of suspensions of BP at every prime and as the map is also rationally injective, we obtain the injectivity of Θ . \square

Note that for any $\lambda \in M\xi_*$ we can express $\Theta(x_\xi \lambda)$ in the form

$$\sum_{i \geq 0} z_i x_H^{i+1} \lambda = \sum_{i \geq 0} z_i \lambda x_H^{i+1},$$

but as the coefficients are non-commutative, we cannot pass λ to the left-hand side, so care has to be taken when calculating in $\text{NSymm}[[x_H]]$.

3. A formal group law over $M\xi_*$

The two evident line bundles η_1, η_2 over $\mathbb{C}P^\infty \times \mathbb{C}P^\infty$ can be tensored together to give a line bundle $\eta_1 \otimes \eta_2$ classified by a map $\mu: \mathbb{C}P^\infty \times \mathbb{C}P^\infty \rightarrow \mathbb{C}P^\infty$ and by naturality we obtain an element $\mu^* x_\xi \in M\xi^2(\mathbb{C}P^\infty \times \mathbb{C}P^\infty)$. We also have

$$(3.1) \quad M\xi^*(\mathbb{C}P^\infty \times \mathbb{C}P^\infty) = \left\{ \sum_{i,j \geq 0} a_{i,j} (x'_\xi)^i (x''_\xi)^j : a_{i,j} \in M\xi_* \right\}$$

as a left $M\xi^* = M\xi_{-*}$ -module, where $x'_\xi, x''_\xi \in M\xi^2(\mathbb{C}P^\infty \times \mathbb{C}P^\infty)$ are obtained by pulling back x_ξ along the two projections. We have

$$\mu^* x_\xi = F_\xi(x'_\xi, x''_\xi) = x'_\xi + x''_\xi + \sum_{i,j \geq 1} a_{i,j} (x'_\xi)^i (x''_\xi)^j,$$

where $a_{i,j} \in M\xi_{2(i+j)-2}$. The notation $F_\xi(x'_\xi, x''_\xi)$ is meant to suggest a power series, but care needs to be taken over the use of such notation. For example, since the tensor product of line bundles is associative up to isomorphism, the formula

$$(3.2a) \quad F_\xi(F_\xi(x'_\xi, x''_\xi), x'''_\xi) = F_\xi(x'_\xi, F_\xi(x''_\xi, x'''_\xi))$$

holds in $M\xi^*(\mathbb{C}P^\infty \times \mathbb{C}P^\infty \times \mathbb{C}P^\infty)$, where $x'_\xi, x''_\xi, x'''_\xi$ denote the pullbacks of x_ξ along the three projections. When considering this formula, we have to bear in mind that the inserted expressions do not commute with each other or coefficients. We also have the identities

$$(3.2b) \quad F_\xi(0, x_\xi) = x_\xi = F_\xi(x_\xi, 0),$$

$$(3.2c) \quad F_\xi(x'_\xi, x''_\xi) = F_\xi(x''_\xi, x'_\xi).$$

Let $\bar{x}_\xi = \gamma^* x_\xi$ denotes the pullback of x_ξ along the map $\gamma: \mathbb{C}P^\infty \rightarrow \mathbb{C}P^\infty$ classifying the inverse $\eta^{-1} = \bar{\eta}$ of the canonical line bundle η . Then $\bar{x}_\xi \in M\xi^2(\mathbb{C}P^\infty)$ and there is a unique expansion

$$\bar{x}_\xi = -x_\xi + \sum_{k \geq 1} c_k x_\xi^{k+1}$$

with $c_k \in M\xi_{2k}$. Since $\eta \otimes \bar{\eta}$ is trivial, this gives the identities

$$F_\xi(x_\xi, \bar{x}_\xi) = 0 = F_\xi(\bar{x}_\xi, x_\xi)$$

and so

$$(3.2d) \quad F_\xi(x_\xi, -x_\xi + \sum_{k \geq 1} c_k x_\xi^{k+1}) = 0 = F_\xi(-x_\xi + \sum_{k \geq 1} c_k x_\xi^{k+1}, x_\xi).$$

To summarize, we obtain the following result.

PROPOSITION 3.1. *The identities (3.2) together show that $F_\xi(x'_\xi, x''_\xi)$ defines a commutative formal group law over the non-commutative ring $M\xi_*$.*

REMARK 3.2. Note however, that most of the classical structure theory for formal group laws over (graded) commutative rings does *not* carry over to the general non-commutative setting. For power series rings over associative rings where the variable commutes with the coefficients most of the theory works as usual. If the variable commutes with the coefficients up to a controlled deviation, then the ring of skew power series still behaves reasonably (see for example [D]), but our case is more general.

4. The splitting of $M\xi$ into wedges of suspensions of BP

In [BR] we showed that there is a splitting of $M\xi$ into a wedge of copies of suspensions of BP locally at each prime p . In the case of MU the inclusion of the bottom summand is given by a map of ring spectra $BP \rightarrow MU_{(p)}$. However, for $M\xi$ this is not the case.

PROPOSITION 4.1. *For each prime p , there is no map of ring spectra $BP \rightarrow M\xi_{(p)}$*

PROOF. We give the proof for an odd prime p , the case $p = 2$ is similar. We set $H_* = (H\mathbb{F}_p)_*$.

Recall that

$$H_*(BP) = \mathbb{F}_p[t_1, t_2, \dots]$$

where $t_r \in H_{2p^r-2}(BP)$ and the \mathcal{A}_* -coaction on these generators is given by

$$\psi(t_n) = \sum_{k=0}^n \zeta_k \otimes t_{n-k}^k,$$

where $\zeta_r \in \mathcal{A}_{2p^r-2}$ is the conjugate of the usual Milnor generator ξ_r , defined in [Ad]. The right action of the Steenrod algebra satisfies

$$\mathcal{P}_*^1 t_1 = -1, \quad \mathcal{P}_*^1 t_2 = -t_1^p, \quad \mathcal{P}_*^p t_2 = 0.$$

Assume that a map of ring spectra $u: BP \rightarrow M\xi_{(p)}$ exists. Then $\mathcal{P}_*^1 u_*(t_1) = u_*(-1) = -1$, hence $w := u_*(t_1) \neq 0$. Notice that

$$\mathcal{P}_*^1(w^{p+1}) = -w^p, \quad \mathcal{P}_*^p(w^{p+1}) = -w.$$

Also, $\mathcal{P}_*^p u_*(t_2) = 0$. This shows that $u_*(t_2)$ cannot be equal to a non-zero multiple of w^{p+1} . Therefore it is not contained in the polynomial subalgebra of $H_*(M\xi_{(p)})$ generated by w^{p+1} and thus it cannot commute with w . This shows that the image of u_* is not a commutative subalgebra of $H_*M\xi_{(p)}$ which contradicts the commutativity of H_*BP . \square

REMARK 4.2. Note that Proposition 4.1 implies that there is no map of ring spectra from MU to $M\xi$, because if such a map existed, we could precompose it p -locally with the ring map $BP \rightarrow MU_{(p)}$ to get a map of ring spectra $BP \rightarrow M\xi_{(p)}$.

5. The real and the quaternionic cases

Analogous to the complex case, the map $\mathbb{R}P^\infty = BO(1) \rightarrow BO$ gives rise to a loop map $\xi_{\mathbb{R}}: \Omega\Sigma\mathbb{R}P^\infty \rightarrow BO$, and hence there is an associated map of associative S -algebras $M\xi_{\mathbb{R}} \rightarrow MO$ on the level of Thom spectra. There is a splitting of MO into copies of suspensions of $H\mathbb{F}_2$. In fact a stronger result holds.

PROPOSITION 5.1. *There is a map of E_2 -spectra $H\mathbb{F}_2 \rightarrow MO$.*

PROOF. The map $\alpha: \mathbb{S}^1 \rightarrow BO$ that detects the generator of the fundamental group of BO gives rise to a double-loop map

$$\Omega^2\Sigma^2\mathbb{S}^1 = \Omega^2\mathbb{S}^3 \rightarrow BO.$$

As the Thom spectrum associated to $\Omega^2\mathbb{S}^3$ is a model of $H\mathbb{F}_2$ by [Mah] with an E_2 -structure [LMSM], the claim follows. \square

Generalizing an argument by Hu-Kriz-May [HKM], Gilmour [G] showed that there is no map of commutative S -algebras $H\mathbb{F}_2 \rightarrow MO$.

The E_2 -structure on the map from Proposition 5.1 cannot be extended to $\xi_{\mathbb{R}}$. On the space level,

$$H_*(\Omega\Sigma\mathbb{R}P^\infty; \mathbb{F}_2) \cong T_{\mathbb{F}_2}(\bar{H}_*(\mathbb{R}P^\infty; \mathbb{F}_2)),$$

where $H_n(\mathbb{R}P^\infty; \mathbb{F}_2)$ is generated by an element x_n .

PROPOSITION 5.2. *There is no map of ring spectra $H\mathbb{F}_2 \rightarrow M\xi_{\mathbb{R}}$.*

PROOF. Assume $\gamma: H\mathbb{F}_2 \rightarrow M\xi_{\mathbb{R}}$ were a map of ring spectra. We consider $\gamma_*: (H\mathbb{F}_2)_*H\mathbb{F}_2 \rightarrow (H\mathbb{F}_2)_*M\xi_{\mathbb{R}}$. Note that $(H\mathbb{F}_2)_*M\xi_{\mathbb{R}}$ is the free associative \mathbb{F}_2 -algebra generated by z_1, z_2, \dots with z_i in degree i being the image of x_i under the Thom-isomorphism.

Under the action of the Steenrod-algebra on $H\mathbb{F}_2$ -homology $Sq_*^1(z_1) = 1$ and hence $Sq_*^1(z_1^3) = z_1^2$ by the derivation property of Sq_*^1 .

In the dual Steenrod algebra we have $Sq_*^1(\xi_1) = 1$ and $Sq_*^2(\xi_2) = \xi_1$ and $Sq_*^1(\xi_2) = 0$.

Combining these facts we obtain

$$(5.1) \quad Sq_*^1(\gamma_*(\xi_1)) = \gamma_*(Sq_*^1\xi_1) = \gamma_*(1) = 1,$$

in particular $\gamma_*(\xi_1) \neq 0$ and thus $\gamma_*(\xi_1) = z_1$.

Similarly,

$$Sq_*^2\gamma_*(\xi_2) = \gamma_*(Sq_*^2\xi_2) = \gamma_*\xi_1 = z_1 \neq 0.$$

The image of γ_* generates a commutative sub- \mathbb{F}_2 -algebra of $(H\mathbb{F}_2)_*M\xi_{\mathbb{R}}$. The only elements in $(H\mathbb{F}_2)_*M\xi_{\mathbb{R}}$ that commute with z_1 are polynomials in z_1 . Assume that $\gamma_*\xi_2 = z_1^3$. Then

$$0 = \gamma_*Sq_*^1\xi_2 = Sq_*^1(z_1^3) = z_1^2 \neq 0,$$

which is impossible. Therefore, $\gamma_*\xi_2$ does not commute with z_1 , so we get a contradiction. \square

Note that Proposition 5.2 implies that there is no loop map $\Omega^2\mathbb{S}^3 \rightarrow \Omega\Sigma\mathbb{R}P^\infty$ that is compatible with the maps to BO since such a map would induce a map of associative S -algebras $H\mathbb{F}_2 \rightarrow M\xi_{\mathbb{R}}$.

A quaternionic model of quasisymmetric functions is given by $H^*(\Omega\Sigma\mathbb{H}P^\infty)$. Here, the algebraic generators are concentrated in degrees that are divisible by 4.

The canonical map $\mathbb{H}P^\infty = BSp(1) \rightarrow BSp$ induces a loop-map $\xi_{\mathbb{H}}: \Omega\Sigma\mathbb{H}P^\infty \rightarrow BSp$ and thus gives rise to a map of associative S -algebras on the level of Thom spectra $M\xi_{\mathbb{H}} \rightarrow MSp$.

Of course, the spectrum MSp is not as well understood as MO and MU . There is a commutative S -algebra structure on MSp [May, pp. 22, 76], but for instance the homotopy groups of MSp are not known in an explicit form.

6. Associative versus commutative orientations

We work with the second desuspension of the suspension spectrum of $\mathbb{C}P^\infty$. Such spectra are inclusion prespectra [EKMM, X.4.1] and thus a map of S -modules from $S = \Sigma^\infty\mathbb{S}^0$ to $\Sigma^{\infty-2}\mathbb{C}P^\infty := \Sigma^{-2}\Sigma^\infty\mathbb{C}P^\infty$ is given by a map from the zeroth space of the sphere spectrum to the zeroth space of $\Sigma^{\infty-2}\mathbb{C}P^\infty$ which in turn is a colimit, namely $\text{colim}_{\mathbb{R}^2 \subset W} \Omega^W \Sigma^{W-\mathbb{R}^2} \mathbb{C}P^\infty$. As a map $\varrho: S \rightarrow \Sigma^{\infty-2}\mathbb{C}P^\infty$ we take the one that is induced by the inclusion $\mathbb{S}^2 = \mathbb{C}P^1 \subset \mathbb{C}P^\infty$.

The commutative S -algebra $S \wedge (\mathbb{N}_0)_+ = S[\mathbb{N}_0]$ has a canonical map $S[\mathbb{N}_0] \rightarrow S$ which is given by the fold map. We can model this via the map of monoids that sends the additive monoid $(\mathbb{N}_0, 0, +)$ to the monoid $(0, 0, +)$; thus $S[\mathbb{N}_0] \rightarrow S$ is a map of commutative S -algebras.

We get a map $S[\mathbb{N}_0] \rightarrow \mathbb{A}(\Sigma^{\infty-2}\mathbb{C}P^\infty)$ by taking the following map on the n th copy of S in $S[\mathbb{N}_0]$. We can view S as $S \wedge \{*\}_+$ where $\{*\}$ is a one-point space. The n -fold space diagonal gives a map

$$\delta_n: S = S \wedge \{*\}_+ \rightarrow S \wedge \underbrace{\{(*, \dots, *)\}}_n \wedge \{*\}_+ \simeq S^{\wedge n}$$

which fixes an equivalence of S with $S^{\wedge n}$. We compose this map with the n -fold smash product of the map $\varrho: S \rightarrow \Sigma^{\infty-2}\mathbb{C}P^\infty$. The maps

$$\varrho^{\wedge n} \circ \delta_n: S \rightarrow (\Sigma^{\infty-2}\mathbb{C}P^\infty)^{\wedge n} \rightarrow \mathbb{A}(\Sigma^{\infty-2}\mathbb{C}P^\infty)$$

together give a map of S -algebras

$$\tau: S[\mathbb{N}_0] \rightarrow \mathbb{A}(\Sigma^{\infty-2}\mathbb{C}P^\infty).$$

Note, however, that $S[\mathbb{N}_0]$ is not central in $\mathbb{A}(\Sigma^{\infty-2}\mathbb{C}P^\infty)$. Thus the coequalizer

$$S \wedge_{S[\mathbb{N}_0]} \mathbb{A}(\Sigma^{\infty-2}\mathbb{C}P^\infty)$$

does not possess any obvious S -algebra structure. Furthermore, there is a natural map

$$S \wedge_{S[\mathbb{N}_0]} \mathbb{A}(\Sigma^{\infty-2}\mathbb{C}P^\infty) \rightarrow M\xi,$$

but this is not a weak equivalence since the $H\mathbb{Z}$ -homology of the left-hand side is the quotient by the left ideal generated by $z_0 - 1$ and thus it is bigger than $H\mathbb{Z}_* M\xi$ which is the quotient by the two-sided ideal generated by $z_0 - 1$.

In the commutative context the pushout of commutative S -algebras is given by the smash product. Hence there is a natural morphism of commutative S -algebras

$$\tilde{\mathbb{P}}(\Sigma^{\infty-2}\mathbb{C}P^\infty) = S \wedge_{\mathbb{P}(S)} \mathbb{P}(\Sigma^{\infty-2}\mathbb{C}P^\infty) \rightarrow MU,$$

where $\tilde{\mathbb{P}}(\Sigma^{\infty-2}\mathbb{C}P^\infty)$ is the pushout in the following diagram of commutative S -algebras

$$\begin{array}{ccc} \mathbb{P}(S) & \longrightarrow & \mathbb{P}(\Sigma^{\infty-2}\mathbb{C}P^\infty) \\ \downarrow & \lrcorner & \downarrow \\ S & \longrightarrow & \tilde{\mathbb{P}}(\Sigma^{\infty-2}\mathbb{C}P^\infty). \end{array}$$

Here, we use the identity map on S to induce the left-hand vertical map of commutative S -algebras and the inclusion of the bottom cell of $\Sigma^{\infty-2}\mathbb{C}P^\infty$ to induce the top map which is a cofibration and therefore $\tilde{\mathbb{P}}(\Sigma^{\infty-2}\mathbb{C}P^\infty)$ is cofibrant. However, the map $\tilde{\mathbb{P}}(\Sigma^{\infty-2}\mathbb{C}P^\infty) \rightarrow MU$ is not a weak equivalence as the next result shows.

LEMMA 6.1. *The canonical map of commutative S -algebras*

$$\tilde{\mathbb{P}}(\Sigma^{\infty-2}\mathbb{C}P^\infty) \rightarrow MU$$

is an equivalence rationally, but not globally. Furthermore, there is a morphism of ring spectra

$$MU \rightarrow \tilde{\mathbb{P}}(\Sigma^{\infty-2}\mathbb{C}P^\infty)$$

which turns MU into a retract of $\tilde{\mathbb{P}}(\Sigma^{\infty-2}\mathbb{C}P^\infty)$.

PROOF. Let \mathbf{k} be a field. The Künneth spectral sequence for the homotopy groups of

$$H\mathbf{k} \wedge (\tilde{\mathbb{P}}(\Sigma^{\infty-2}\mathbb{C}P^\infty)) \simeq H\mathbf{k} \wedge_{\mathbb{P}_{H\mathbf{k}}(H\mathbf{k})} \mathbb{P}_{H\mathbf{k}}(\Sigma^{-2}H\mathbf{k} \wedge \mathbb{C}P^\infty)$$

has E^2 -term

$$E_{*,*}^2 = \mathrm{Tor}_{*,*}^{\pi_*(\mathbb{P}_{H\mathbf{k}}(H\mathbf{k}))}(\mathbf{k}, \pi_*(\mathbb{P}_{H\mathbf{k}}(\Sigma^{-2}H\mathbf{k} \wedge \mathbb{C}P^\infty))).$$

When $\mathbf{k} = \mathbb{Q}$, $\pi_*(\mathbb{P}_{H\mathbb{Q}}(H\mathbb{Q}))$ is a polynomial algebra on a zero-dimensional class x_0 and

$$(6.1) \quad \pi_*(\mathbb{P}_{H\mathbb{Q}}(\Sigma^{-2}H\mathbb{Q} \wedge \mathbb{C}P^\infty)) \cong \mathbb{Q}[x_0, x_1, \dots],$$

where $|x_i| = 2i$. Thus

$$\pi_*(H\mathbb{Q} \wedge (\tilde{\mathbb{P}}(\Sigma^{\infty-2}\mathbb{C}P^\infty))) \cong \mathbb{Q}[x_1, x_2, \dots] \cong H\mathbb{Q}_*(MU).$$

However, when $\mathbf{k} = \mathbb{F}_p$ for a prime p , the freeness of the commutative S -algebras $\mathbb{P}(S)$ and $\mathbb{P}(\Sigma^{\infty-2}\mathbb{C}P^\infty)$ implies that $(H\mathbb{F}_p)_*(\mathbb{P}(\Sigma^{\infty-2}\mathbb{C}P^\infty))$ is a free $(H\mathbb{F}_p)_*(\mathbb{P}(S))$ -module and thus the E^2 -term reduces to the tensor product in homological degree zero. Note that this tensor product contains elements of odd degree, but $(H\mathbb{F}_p)_*(MU)$ doesn't.

Using the orientation for line bundles given by the canonical inclusion

$$\Sigma^{\infty-2}\mathbb{C}P^\infty \rightarrow \tilde{\mathbb{P}}(\Sigma^{\infty-2}\mathbb{C}P^\infty),$$

we have a map of ring spectra

$$\varphi: MU \rightarrow \tilde{\mathbb{P}}(\Sigma^{\infty-2}\mathbb{C}P^\infty).$$

The inclusion map $\mathbb{C}P^\infty = BU(1) \rightarrow BU$ gives rise to the canonical map $\sigma: \Sigma^{\infty-2}\mathbb{C}P^\infty \rightarrow MU$ and with this orientation we get a morphism of commutative S -algebras

$$\theta: \tilde{\mathbb{P}}(\Sigma^{\infty-2}\mathbb{C}P^\infty) \rightarrow MU,$$

such that the composite $\theta \circ \varphi \circ \sigma$ agrees with σ , hence $\theta \circ \varphi$ is homotopic to the identity on MU . \square

Using topological André-Quillen homology, $TAQ_*(-)$, we can show that the map of ring spectra φ cannot be rigidified to a map $\tilde{\varphi}$ of commutative S -algebras in such a way that the composite $\theta \circ \tilde{\varphi}$ is a weak-equivalence. By Basterra-Mandell [BM],

$$TAQ_*(MU|S; H\mathbb{F}_p) \cong (H\mathbb{F}_p)_*(\Sigma^2 ku),$$

while [BGR, proposition 1.6] together with subsequent work of the first named author [Ba] gives

$$TAQ_*(\tilde{\mathbb{P}}(\Sigma^{\infty-2}\mathbb{C}P^\infty)|S; H\mathbb{F}_p) \cong (H\mathbb{F}_p)_*(\Sigma^{\infty-2}\mathbb{C}P_2^\infty),$$

where $\mathbb{C}P_2^\infty = \mathbb{C}P^\infty/\mathbb{C}P^1$ is the cofiber of the inclusion of the bottom cell.

PROPOSITION 6.2. *For a prime p , there can be no morphism of commutative $S_{(p)}$ -algebras*

$$\theta_{(p)}: MU_{(p)} \rightarrow (\tilde{\mathbb{P}}\Sigma^{\infty-2}\mathbb{C}P^\infty)_{(p)}$$

for which $\sigma_{(p)} \circ \theta_{(p)}$ is a weak equivalence. Hence there can be no morphism of commutative S -algebras

$$\theta: MU \rightarrow \tilde{\mathbb{P}}\Sigma^{\infty-2}\mathbb{C}P^\infty$$

for which $\sigma \circ \theta$ is a weak equivalence.

PROOF. It suffices to prove the first result, and we will assume that all spectra are localised at p . Assume such a morphism θ existed. Then by naturality of the functor of Kähler differentials, Ω_S , there are (derived) morphisms of MU -modules and a commutative diagram

$$\begin{array}{ccccc} & & \sim & & \\ & \searrow & & \nearrow & \\ \Omega_S(MU) & \xrightarrow{\theta_*} & \Omega_S(\tilde{\mathbb{P}}\Sigma^{\infty-2}\mathbb{C}P^\infty) & \xrightarrow{\sigma_*} & \Omega_S(MU) \end{array}$$

which by [BM] induce a commutative diagram in $TAQ_*(-; H\mathbb{F}_p)$ of the following form:

$$\begin{array}{ccccc} & & \cong & & \\ & \searrow & & \nearrow & \\ H_*(\Sigma^2 ku; \mathbb{F}_p) & \xrightarrow{\theta_*} & H_*(\Sigma^{\infty-2}\mathbb{C}P_2^\infty; \mathbb{F}_p) & \xrightarrow{\sigma_*} & H_*(\Sigma^2 ku; \mathbb{F}_p). \end{array}$$

It is standard that

$$H_n(\Sigma^{\infty-2}\mathbb{C}P_2^\infty; \mathbb{F}_p) = \begin{cases} \mathbb{F}_p & \text{if } n \geq 2 \text{ and is even,} \\ 0 & \text{otherwise.} \end{cases}$$

On the other hand, when $p = 2$,

$$H_*(ku; \mathbb{F}_2) = \mathbb{F}_2[\zeta_1^2, \zeta_2^2, \zeta_3, \zeta_4, \dots] \subseteq \mathcal{A}(2)_*$$

with $|\zeta_s| = 2^s - 1$, while when p is odd

$$\Sigma^2 ku_{(p)} \sim \bigvee_{1 \leq r \leq p-1} \Sigma^{2r} \ell,$$

where ℓ is the Adams summand with

$$H_*(\ell; \mathbb{F}_2) = \mathbb{F}_2[\zeta_1, \zeta_2, \zeta_3, \dots] \otimes \Lambda(\bar{\tau}_r : r \geq 2),$$

for $|\zeta_s| = 2p^s - 2$ and $|\bar{\tau}_s| = 2p^s - 1$. Hence no such θ can exist. \square

PROPOSITION 6.3. *There are commutative S -algebras E which possess a map of commutative S -algebras*

$$\tilde{\mathbb{P}}(\Sigma^{\infty-2}\mathbb{C}P^{\infty}) \rightarrow E$$

that cannot be extended to a map of commutative S -algebras $MU \rightarrow E$.

PROOF. Matthew Ando [An] constructed complex orientations for the Lubin-Tate spectra E_n which are H_{∞} -maps $MU \rightarrow E_n$. However, in [JN], Niles Johnson and Justin Noël showed that none of these are p -typical for all primes up to at least 13 (and subsequently verified for primes up to 61). For any p -typical orientation there is a map of ring spectra $MU \rightarrow E_n$, but this map cannot be an H_{∞} -map and therefore is not a map of commutative S -algebras. \square

Acknowledgements

The authors would like to thank Jack Morava and Nitu Kitchloo for encouraging them to work out details for some of the loose ends left over from our earlier paper. The authors also thank Tyler Lawson for introducing the first named author to the spectrum $\tilde{\mathbb{P}}(\Sigma^{\infty-2}\mathbb{C}P^{\infty})$ and its universal property.

References

- [Ad] J. F. Adams, *Stable homotopy and generalised homology*, University of Chicago Press, Chicago, Ill., 1974. Chicago Lectures in Mathematics. MR0402720 (53 #6534)
- [An] Matthew Ando, *Isogenies of formal group laws and power operations in the cohomology theories E_n* , Duke Math. J. **79** (1995), no. 2, 423–485, DOI 10.1215/S0012-7094-95-07911-3. MR1344767 (97a:55006)
- [Ba] Andrew Baker, *Calculating with topological André-Quillen theory, I: Homotopical properties of universal derivations and free commutative S -algebras*, arXiv:1208.1868.
- [BGR] Andrew Baker, Helen Gilmour, and Philipp Reinhard, *Topological André-Quillen homology for cellular commutative S -algebras*, Abh. Math. Semin. Univ. Hambg. **78** (2008), no. 1, 27–50, DOI 10.1007/s12188-008-0005-9. MR2501527 (2010c:55006)
- [BR] Andrew Baker and Birgit Richter, *Quasisymmetric functions from a topological point of view*, Math. Scand. **103** (2008), no. 2, 208–242. MR2484353 (2009k:55025)
- [BM] Maria Basterra and Michael A. Mandell, *Homology and cohomology of E_{∞} ring spectra*, Math. Z. **249** (2005), no. 4, 903–944, DOI 10.1007/s00209-004-0744-y. MR2126222 (2005m:55016)
- [D] François Dumas, *Skew power series rings with general commutation formula*, Theoret. Comput. Sci. **98** (1992), no. 1, 99–114, DOI 10.1016/0304-3975(92)90382-P. Second Workshop on Algebraic and Computer-theoretic Aspects of Formal Power Series (Paris, 1990). MR1169128 (93h:16050)
- [EKMM] A. D. Elmendorf, I. Kriz, M. A. Mandell, and J. P. May, *Rings, modules, and algebras in stable homotopy theory*, Mathematical Surveys and Monographs, vol. 47, American Mathematical Society, Providence, RI, 1997. With an appendix by M. Cole. MR1417719 (97h:55006)
- [G] Helen Gilmour, *Nuclear and Minimal Atomic S -algebras*, PhD Thesis, University of Glasgow (2006).
- [HKM] P. Hu, I. Kriz, and J. P. May, *Cores of spaces, spectra, and E_{∞} ring spectra*, Homology Homotopy Appl. **3** (2001), no. 2, 341–354. Equivariant stable homotopy theory and related areas (Stanford, CA, 2000). MR1856030 (2002j:55005)
- [JN] Niles Johnson and Justin Noel, *For complex orientations preserving power operations, p -typicality is atypical*, Topology Appl. **157** (2010), no. 14, 2271–2288, DOI 10.1016/j.topol.2010.06.007. MR2670503 (2011g:55009)
- [LMSM] L. G. Lewis Jr., J. P. May, M. Steinberger, and J. E. McClure, *Equivariant stable homotopy theory*, Lecture Notes in Mathematics, vol. 1213, Springer-Verlag, Berlin, 1986. With contributions by J. E. McClure. MR866482 (88e:55002)

- [Mah] Mark Mahowald, *Ring spectra which are Thom complexes*, Duke Math. J. **46** (1979), no. 3, 549–559. MR544245 (81f:55010)
- [May] J. Peter May, *E_∞ ring spaces and E_∞ ring spectra*, Lecture Notes in Mathematics, Vol. 577, Springer-Verlag, Berlin, 1977. With contributions by Frank Quinn, Nigel Ray, and Jørgen Tornehave. MR0494077 (58 #13008)
- [MK] Jack Morava & Nitu Kitchloo, *The Baker-Richter spectrum as cobordism of quasitoric manifolds*, [arxiv:1201.3127](https://arxiv.org/abs/1201.3127).
- [Sch] Christian Schlichtkrull, *Thom spectra that are symmetric spectra*, Doc. Math. **14** (2009), 699–748. MR2578805 (2011f:55017)

SCHOOL OF MATHEMATICS & STATISTICS, UNIVERSITY OF GLASGOW, GLASGOW G12 8QW, SCOTLAND.

E-mail address: a.baker@maths.gla.ac.uk

URL: <http://www.maths.gla.ac.uk/~ajb>

FACHBEREICH MATHEMATIK DER UNIVERSITÄT HAMBURG, 20146 HAMBURG, GERMANY.

E-mail address: richter@math.uni-hamburg.de

URL: <http://www.math.uni-hamburg.de/home/richter/>