

# **Infinite-dimensional Lie Theory for Gauge Groups**

Christoph Wockel  
Fachbereich Mathematik  
Technische Universität Darmstadt

[www.mathematik.tu-darmstadt.de/~wockel](http://www.mathematik.tu-darmstadt.de/~wockel)

Nancy, June 10, 2005

# **Overview**

1. Introduction and Motivation
2. The Topology of Gauge Groups
3. Central Extensions of Gauge Groups

# 1. Introduction and Motivation

**Setting:**

$$P \xrightarrow{\pi} M$$

smooth  $K$ -principal bundle (possibly infinite-dimensional), i.e.

- $P \times K \rightarrow P$  smooth free action
- $P/K \cong M$  and  $\pi(x) = x.K$
- $\pi$  admits smooth local sections

**Examples:**

- trivial bundles  $M \times K \xrightarrow{\text{pr}} M$
- homogeneous spaces  $G \xrightarrow{q} G/H$

**Physical Motivation:**

$M \leftrightarrow$  space time

$\text{Diff}(M) \leftrightarrow$  symmetry group of  $M$

$\text{Conn}(P) \leftrightarrow$  potentials of forces

$\text{Gau}(P) \leftrightarrow$  symmetry group of  $\text{Conn}(P)$

## Automorphisms:

$$\text{Aut}(P) := \{f \in \text{Diff}(P) : f(p \cdot k) = f(p) \cdot k\}$$

$$\begin{aligned} \text{Gau}(P) &:= \{f \in \text{Aut}(P) : f(p) = p \cdot \gamma_f(p) \\ &\quad \gamma_f \text{ smooth, } \gamma_f(p \cdot k) = \gamma_f(p) \cdot k\} \end{aligned}$$

$$\Rightarrow \text{Gau}(P) \cong C^\infty(P, K)^K$$

## Aim:

- Topologise  $C^\infty(P, K)^K$
- Calculate  $\pi_n(C^\infty(P, K)^K)$
- Construct central extensions of  $C^\infty(P, K)^K$

## Connection to mapping groups:

- $P$  trivial (i.e.  $\sigma : M \rightarrow P$  section)  
 $\Rightarrow \sigma^* : C^\infty(P, K)^K \rightarrow C^\infty(M, K)$  iso
- $K$  abelian &  $f \in C^\infty(P, K)^K$   
 $\Rightarrow f(p \cdot k) = k^{-1} \cdot f(p) \cdot k = f(p)$   
 $\Rightarrow f^\# : P/K \cong M \rightarrow K$

## 2. The Topology of Gauge Groups

From now on :  $\text{Gau}(P) = C^\infty(P, K)^K$

**Definition:**  $f \in C^\infty(P, K)$

$$\Rightarrow T^n f \in C(T^n P, T^n K)$$

$$\Rightarrow C^\infty(P, K)^K \hookrightarrow \prod_{n=0}^{\infty} C(T^n P, T^n K)_c$$

leads to group topology on  $C^\infty(P, K)^K$

**Theorem** (W.,'03):  $M$  compact,  
 $\exp : \mathfrak{k} \rightarrow K$  restricts to diffeomorphism

$\Rightarrow C^\infty(P, K)^K$  is  $\infty$ -dimensional Lie group,  
 $\exp_* : C^\infty(P, \mathfrak{k})^K \rightarrow C^\infty(P, K)^K$

**Lie algebra:**  $C^\infty(P, \mathfrak{k})^K \cong \text{sec}(P \times_{\text{Ad}} \mathfrak{k})$

From now on assume  $M$  compact,  $K$   
locally exponential!

## **Homotopy groups:**

**Aim:** Calculation of  $\pi_n(C^\infty(P, K)^K)$

**Theorem** (W.,'04):

$$\iota : C^\infty(P, K)^K \rightarrow C(P, K)^K$$

induces isomorphisms

$$\pi_n(\iota) : \pi_n(C^\infty(P, K)^K) \rightarrow \pi_n(C(P, K)^K)$$

$\Rightarrow$  suffices to consider  $C(P, K)^K$

Consider bundles over orientable compact surfaces with connected  $K$ .

**Fact:**  $\partial M \neq \emptyset \Rightarrow P \cong M \times K$

$$\Rightarrow C(P, K)^K \cong C(M, K)$$

**Theorem** (W.,'05): If  $K = K_0$ ,  $\partial M \neq \emptyset$ , then

$$\pi_n(C(P, K)^K) \cong \pi_{n+1}(K)^{2g-m-1} \oplus \pi_n(K)$$

with  $g = \text{gen}(M)$  and  $m = \#\text{comp}(\partial M)$ .

$$\partial M = \emptyset \Rightarrow \text{Bun}(K, M) \cong \pi_1(K)$$

$\Rightarrow P$  determined by  $\gamma \in C_*(\mathbb{S}^1, K)$

$$\begin{aligned} &\Rightarrow_{\text{torus}} C(P, K)^K \cong \\ &\quad \{f \in C([0, 1]^2, K) : f(0, s) = f(1, s) \\ &\quad \quad f(t, 0) = \gamma(t)^{-1} \cdot f(t, 1) \cdot \gamma(t)\} \end{aligned}$$

$$\begin{aligned} &\Rightarrow C(P, K)_*^K \cong \{f : f(0) = e\} \cong \\ &\quad C_*(\mathbb{S}^2, K) \times C_*(\mathbb{S}^1, K)^{2g} \end{aligned}$$

**Theorem** (W., '05): If  $K = K_0$ ,  $\partial M = \emptyset$ ,  $g = \text{gen}(M)$ , then

$$\pi_n(C(P, K)_*^K) \cong \pi_{n+2}(K) \oplus \pi_{n+1}(K)^{2g}.$$

**Problem:**  $C(P, K)_*^K \rightarrow C(P, K)^K \rightarrow K$   
does not split canonically!

$\Rightarrow$  Obtain an exact homotopy sequence

$$\begin{aligned} \dots &\rightarrow \pi_{n+1}(K) \rightarrow \pi_n(C(P, K)_*^K) \\ &\quad \rightarrow \pi_n(C(P, K)^K) \rightarrow \pi_n(K) \rightarrow \dots \end{aligned}$$

### 3. Central Extensions of Gauge Groups (according to Losev et.al.)

- $G := C^\infty(P, K)_0^K$
- $\mathfrak{g} := C^\infty(P, \mathfrak{k})^K \cong \sec(P \times_{\text{Ad}} \mathfrak{k})$

**Aim:** Construct central extensions

$$Z \hookrightarrow \widehat{G} \twoheadrightarrow G$$

**Philosophy:**

- Central extensions are parametrised by cohomology classes  $H^2(\mathfrak{g}, \mathfrak{z})$ ,  $H^2(G, Z)$ .
- Construct cocycle in  $H^2(\mathfrak{g}, \mathfrak{z})$ , then try to 'integrate'.

**Mapping Groups** (Pressley&Segal,86;  
Maier&Neeb,03): Construction of central  
extensions for  $C^\infty(M, K)_0$  for  $\dim(K) < \infty$ .

## Central extensions of $\mathfrak{g}$ : Construct cocycle

$$\omega_{\kappa, \nabla} : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{z}$$

- $\kappa : \mathfrak{k} \times \mathfrak{k} \rightarrow V$  invariant, symmetric, bilinear
- $\nabla : \mathfrak{g} \times \mathcal{V}(M) \rightarrow \mathfrak{g}$ ,  $(\xi, X) \mapsto \nabla_\xi X$  connection (or covariant derivative)
- $\mathfrak{z} := \Omega^1(M, V)/dC^\infty(M, V) \supseteq H_{dR}^1(M, V)$

$$\omega_{\kappa, \nabla}(\xi, \eta) := [\kappa(\xi, \nabla\eta)] \in \mathfrak{z}$$

$\nabla, \nabla'$  connect.  $\Rightarrow \omega_{\kappa, \nabla} - \omega_{\kappa, \nabla'}$  coboundary  
 $\Rightarrow [\omega_\kappa] := [\omega_{\kappa, \nabla}] \in H^2(\mathfrak{g}, \mathfrak{z})$  independent of choice of  $\nabla$ , yields central extension

$$\mathfrak{z} \hookrightarrow \underbrace{\mathfrak{z} \oplus_{\omega_\kappa} \mathfrak{g}}_{\widehat{\mathfrak{g}}} \twoheadrightarrow \mathfrak{g}.$$

**Central extension of  $G$ :** Period map

$$\text{per}_\omega : \pi_2(G) \times \pi_1(G) \rightarrow \mathfrak{z} \times \text{Lin}(\mathfrak{z}, \mathfrak{g})$$

$\text{im}(\text{per}_\omega) = \Gamma \times 0 \Rightarrow$  ex. corresponding central extension

$$Z \hookrightarrow \widehat{G} \twoheadrightarrow G$$

**Reduction Theorem** (W.,'03): If  $K = K_0$ , then

$$\text{im}(\text{per}_\omega) = \Gamma \times 0 \Leftrightarrow \text{im}(\text{per}_{\omega, \mathbb{S}^1}) = \Gamma_{\mathbb{S}^1} \times 0$$

Is in particular satisfied if  $\dim(K) < \infty$

**Universality?** For mapping groups, finite-dim. s.-s.  $\mathfrak{k}$ : relies on  $\mathfrak{k} \leq C^\infty(M, \mathfrak{k})$ , i.e.

$$C_*^\infty(M, \mathfrak{k}) \hookrightarrow C^\infty(M, \mathfrak{k}) \twoheadrightarrow \mathfrak{k}$$

splits. For gauge algebras not true, same problem as before!

## Conclusion

- $C^\infty(P, K)^K$  is  $\infty$ -dim. Lie group
- topology on  $C^\infty(P, K)^K$  accessible
- can construct central extensions
- problem:  $\mathfrak{k}$  no canonical subalgebra of  $C^\infty(P, \mathfrak{k})^K$