

ON FIXING BOUNDARY POINTS OF TRANSITIVE HYPERBOLIC GRAPHS

AGELOS GEORGAKOPOULOS AND MATTHIAS HAMANN

ABSTRACT. We show that there is no one-ended, locally finite, planar, hyperbolic graph such that the stabilizer of one of its hyperbolic boundary points acts transitively on the vertices of the graph. This gives a partial answer to a question by Kaimanovich and Woess.

1. INTRODUCTION

In [13], Woess asked for a classification of the multi-ended locally finite graphs such that a subgroup of their automorphism group acts transitively on the vertices and fixes an end. This problem was solved by Möller [11] by showing that these graphs are quasi-isometric to *semi-regular* trees, that means that all vertices in each set of its natural bipartition have the same degree. For one-ended graphs the above question makes no sense, however it becomes interesting if one refines the ends by considering some other boundary. Kaimanovich and Woess [10] considered this question with respect to the Gromov-hyperbolic boundary.

As the hyperbolic boundary is a refinement of the space of ends of a graph, an end that contains a hyperbolic boundary point fixed by a subgroup of the automorphism group of the graph is also fixed by that group. Thus, the situation is solved in the cases where the hyperbolic graph has infinitely many ends by Möller's aforementioned result. As infinite locally finite transitive graphs have either 1, 2, or infinitely many ends (compare with [6, Corollary 4]) and the two-ended graphs are quasi-isometric to \mathbb{R} , the only case that remains to be discussed is the one-ended. Thus Kaimanovich and Woess [10] asked:

Question 1. [10, Section 6.4] *Does there exist a one-ended locally finite hyperbolic graph G and a group acting transitively on VG and fixing precisely one hyperbolic boundary point?*

Perhaps the only known result regarding Question 1 is that, as proved in [7, Théorème 8.30] and in [14, Section 4.D], compare also with [8], for a finitely generated hyperbolic group G with infinite boundary, the action of G on any of its locally finite Cayley graphs fixes no hyperbolic boundary point.

The main result of this paper is that no graph satisfying the assertion of Question 1 has an embedding into the Euclidean plane. As an intermediate step, we obtain a general result (Lemma 3.3) proving the existence of certain types of automorphisms in a group as in Question 1 that might help prove the general case.

1991 *Mathematics Subject Classification.* 05E18, 20F67.

Key words and phrases. hyperbolic groups, group action, boundary, planar graphs.

Both authors were supported by FWF grant P-19115-N18.

2. DEFINITIONS AND BASIC FACTS

2.1. Hyperbolic graphs. In this section we define hyperbolic graphs and various related objects. For a more detailed introduction to hyperbolicity, we refer to [1, 4, 7, 8] and [15, Chapter 22]. We will use the terminology of [5].

Let $G = (VG, EG)$ be a graph. A *geodesic* is a path between two vertices x and y with length $d(x, y)$, i.e. the $x - y$ distance in the graph. The graph G is called δ -*hyperbolic* for a $\delta \geq 0$ if it is locally finite¹ and if for every three vertices $x, y, z \in VG$, for every choice of three geodesics $\pi_{xy}, \pi_{yz}, \pi_{zx}$ joining x, y, z in pairs, and for any point ξ on π_{xy} , there is a point on π_{yz} or π_{zx} having distance at most δ to ξ . Note that ξ might be a vertex or an inner point of an edge². We call G *hyperbolic* if there exists a $\delta \geq 0$ such that G is δ -hyperbolic.

A *ray* is a one-way infinite path. Two rays are *equivalent* if for any finite set S of vertices they both lie eventually in the same component of $G - S$. The equivalence classes of this relation are the *ends* of G .

A (bi-)infinite path is called a *(bi-)infinite geodesic* if every finite subpath of it is a geodesic. Two infinite geodesics $\pi = x_1x_2\dots$ and $\pi' = y_1y_2\dots$ are *equivalent* if they lie within bounded distance from each other. It is well-known, see for example [15, (22.12)], that this defines an equivalence relation. A *hyperbolic boundary point* is an equivalence class of infinite geodesics and the *hyperbolic boundary* ∂G is the set of hyperbolic boundary points. Let \widehat{G} denote $G \cup \partial G$.

By [7, Proposition 7.2.9], we can equip \widehat{G} with a topology such that it is a compact space and such that every infinite geodesic converges to the hyperbolic boundary point it is contained in.

Proposition 2.1. [15, (22.11) and (22.15)] *Let G be a hyperbolic graph with two distinct boundary points η and ν . Let o be a vertex in G , $x_1x_2\dots$ an infinite geodesic converging to η , and $y_1y_2\dots$ an infinite geodesic converging to ν . Then the following two properties hold:*

- (i) *There is an infinite geodesic in G starting at o and having only finitely many vertices outside $\{x_i \mid i \in \mathbb{N}\}$.*
- (ii) *There is a bi-infinite geodesic D having only finitely many vertices outside $\{x_i \mid i \in \mathbb{N}\} \cup \{y_i \mid i \in \mathbb{N}\}$. One side of D converges to η , the other to ν . \square*

Equivalent infinite geodesics stay close to each other:

Proposition 2.2. [15, (22.12)] *If $x_1x_2\dots$ and $y_1y_2\dots$ are equivalent infinite geodesics in a hyperbolic graph, then there is a $k \in \mathbb{Z}$ with $d(x_n, y_{n-k}) \leq 2\delta$ for all but finitely many n . \square*

Let $\gamma > 1, c \geq 0$. An infinite path $x_0x_1\dots$ in G is called an *infinite (γ, c) -quasi-geodesic* if $d(x_i, x_j) \leq \gamma|i - j| + c$ for all $i, j \in \mathbb{N}$. A bi-infinite (γ, c) -quasi-geodesic is defined similarly. Hence an infinite geodesic is a $(1, 0)$ -quasi-geodesic. If the constants γ, c are not important then we just speak of *quasi-geodesics*.

The next proposition shows, that in every hyperbolic graph the geodesics and quasi-geodesics lie close to each other, see also [1, Proposition 3.3], [4, 3.1.3], [7, Théorème 5.6, Théorème 5.11], and [8, 7.2.A].

¹The requirement of local-finiteness can be dropped from the definition of hyperbolic, but as all graphs in this paper are locally finite we keep it for simplicity.

²We are considering the graphs as 1-simplices, which means that every edge is assumed to be an isometric image of the unit interval $[0, 1]$.

Proposition 2.3. [4, Théorème 3.1.4] *Let G be a δ -hyperbolic graph. For all $\gamma_1 \geq 1, \gamma_2 \geq 0$ there is a constant $\kappa = \kappa(\delta, \gamma_1, \gamma_2)$ such that for every two vertices $x, y \in VG$ every (γ_1, γ_2) -quasi-geodesic between x and y lies in a κ -neighborhood around every geodesic between x and y and vice versa.*

Furthermore, this extends to (bi-)infinite (γ_1, γ_2) -quasi-geodesics as well as (bi-)infinite geodesics. \square

The following result is [1, Proposition 3.2] (see also [8, 8.1.D] and [7, Proposition 8.21]).

Proposition 2.4. *Let G be a transitive δ -hyperbolic graph. Let $x \in VG$ and $\alpha \in \text{Aut}(G)$ be such that the orbit of x under α is infinite. Then the set $\{\dots, x\alpha^{-1}, x, x\alpha, \dots\}$ lies on a bi-infinite (κ, λ) -quasi-geodesic for constants $\kappa \geq 1, \lambda \geq 0$ that depend only on δ and $d(x, x\alpha)$.* \square

In terms of Section 3, the automorphism α in Proposition 2.4 is a hyperbolic element; it fixes the two boundary points to which the two sets $\{x\alpha^i \mid i \in \mathbb{N}\}$ and $\{x\alpha^{-i} \mid i \in \mathbb{N}\}$ converge.

2.2. Planar graphs. A graph is *planar* if it admits an embedding, as a 1-complex, into the Euclidean plane. Such embeddings are called *planar* embeddings. A *face* of a planar embedding is a component of the complement of its image, that is, a maximal connected subset of the plane to which no vertex or edge is mapped.

An embedding of G is called *consistent* if, intuitively, it embeds every vertex in a similar way in the sense that the group action carries faces to faces. Let us make this more precise. Given an embedding σ of a graph G , we consider for every vertex x the embedding of the edges incident with x , and define the *spin* of x to be the cyclic order of the set $\{xy \mid y \in N(x)\}$ in which xy_1 is a successor of xy_2 whenever the edge xy_2 comes immediately after the edge xy_1 as we move clockwise around x .

Call an automorphism α of G *spin-preserving* if for every $x \in VG$ the spin of $x\alpha$ is the image of the spin of x in σ . Call it *spin-reversing* if for every $x \in VG$ the spin of $x\alpha$ is the reverse of the image of the spin of x in σ . Call an automorphism *consistent* if it is spin-preserving or spin-reversing in σ . Finally, call the embedding σ *consistent* if every automorphism of G is consistent in σ .

Given a planar embedding of a graph G , we define a *facial path* to be a path of G contained in the closure of a face. It is straightforward to check that σ is consistent if and only if every automorphism of G maps every facial path to a facial path. Thus the following classical result, proved by Whitney [12, Theorem 11] for finite graphs and by Imrich [9] for infinite ones, implies that all planar embeddings of a 3-connected transitive graph (defined below) are consistent.

Theorem 2.5. *Let G be a 3-connected graph embedded in the sphere. Then every automorphism of G maps each facial path to a facial path. Thus every automorphism of G is consistent.* \square

The *connectivity* of a graph is the cardinality of a smallest vertex set whose deletion disconnects the graph. A graph is 3-connected if its connectivity is at least 3. The next result is due to Babai and Watkins [3], see also [2, Lemma 2.4].

Lemma 2.6. [3, Theorem 1] *Let G be a locally finite connected transitive graph that has precisely one end. Let d be the degree of any of its vertices. Then the connectivity of G is at least $3(d+1)/4$.* \square

We deduce from Lemma 2.6 and Theorem 2.5 that every transitive planar graph with precisely one end has a consistent embedding in the Euclidean plane. This means that for every transitive planar one-ended graph G there are only two possibilities for the spin of each vertex, obtained from each other by reversing the order.

3. PROOF OF THE MAIN THEOREM

We shall prove that every planar hyperbolic graph answers Question 1 in the negative. Before we directly attack the question in the situation of planar graphs, we prove a general lemma (Lemma 3.3) which might help to give a negative answer to the question in the general case.

Let us recall the notions of elliptic and hyperbolic automorphisms. Let G be a hyperbolic graph.

- (i) An automorphism of G is called *elliptic* if it fixes a finite set of vertices.
- (ii) An automorphism α of G is called *hyperbolic* if it is not elliptic and fixes precisely two boundary points.

Remark that for every $x \in VG$ and for any hyperbolic element α by the aforementioned results in Section 2.1, the sequence $(x\alpha^n)_{n \in \mathbb{N}}$ converges to one of the boundary points fixed by α and $(x\alpha^{-n})_{n \in \mathbb{N}}$ converges to the other fixed one. Furthermore, we call the boundary point to which the sequence $(x\alpha^n)_{n \in \mathbb{N}}$ converges the *direction* of α .³

For automorphism groups of hyperbolic graphs, there is the following classification of their elements; compare with [4, Chapitre 9].

Lemma 3.1. *Any automorphism of a hyperbolic graph is either elliptic or hyperbolic.* \square

A lemma that ensures an automorphism to be a hyperbolic element is the following which can be found in the book [7] of Ghys and de la Harpe.

Lemma 3.2. [7, Corollaire 8.22] *Let G be a δ -hyperbolic graph and α an automorphism of G . If $d(x, x\alpha^2) > d(x, x\alpha) + 18\delta$, for an $x \in VG$, then α is a hyperbolic element.* \square

We now show the existence of certain elliptic and hyperbolic elements.

Lemma 3.3. *Let G be a one-ended δ -hyperbolic graph and Γ be a group acting transitively on G such that Γ fixes a hyperbolic boundary point ω of G . Then the following statements hold.*

- (i) *For every two vertices $x, y \in VG$ with $d(x, y) > 20\delta$ that lie on a common bi-infinite geodesic between ω and another hyperbolic boundary point, there exists a hyperbolic element h in Γ with $xh = y$.*
- (ii) *The stabilizer of any vertex is non-trivial.*
- (iii) *In the stabilizer of any vertex, there are two non-trivial elements whose product is also non-trivial and contained in that stabilizer.*

Proof. To prove (i) let x, y lie on a common bi-infinite geodesic π as in the assertion with $d(x, y) > 20\delta$ such that x separates y from ω on π . Let $\alpha \in \Gamma$ with $x\alpha = y$.

³Sometimes, this is also called *attractor* or *attracting point*.

Due to the hyperbolicity of G , there is a vertex z on $y\pi$ that has distance at most δ to $y\alpha$. Then we have

$$d(y, z) + \delta \geq d(y, z) + d(z, y\alpha) \geq d(y, y\alpha) > 20\delta.$$

This implies

$$\begin{aligned} d(x, x\alpha^2) + \delta &\geq d(x, x\alpha^2) + d(x\alpha^2, z) \geq d(x, z) \\ &= d(x, x\alpha) + d(x\alpha, z) > d(x, x\alpha) + 19\delta \end{aligned}$$

and the claim follows by Lemma 3.2.

For the proof of (ii), let α_0 be a hyperbolic element in Γ . Then α_0 fixes ω and precisely one further boundary point η_0 . We assume that the direction of α_0 is η_0 . For any $x_0 \in VG$, there are constants $c_1 \geq 1, c_2 \geq 0$ such that the vertices $x_0\alpha_0^i$, $i \in \mathbb{Z}$, lie on a bi-infinite (c_1, c_2) -quasi-geodesic π_0 by Proposition 2.4. Note that c_1 and c_2 depend only on δ and $d(x_0, x_0\alpha_0)$.

We are now going to construct a sequence $(x_i)_{i \in \mathbb{N}}$ of vertices in G , a sequence $(\pi_i)_{i \in \mathbb{N}}$ of bi-infinite (c_1, c_2) -quasi-geodesics, a sequence $(\eta_i)_{i \in \mathbb{N}}$ of hyperbolic boundary points, a sequence $(\alpha_i)_{i \in \mathbb{N}}$ of hyperbolic elements of Γ , and a sequence $(\beta_i)_{i \in \mathbb{N}}$ of automorphisms of G , with the following properties:

- (1) the orbit of x_i under α_i lies on π_i ;
- (2) any infinite geodesic contained in π_i converges to either ω or η_i ;
- (3) η_i is the direction of α_i , and
- (4) x_i has distance more than 2κ to all π_j with $j < i$.

For this, let $\kappa = \kappa(\delta, c_1, c_2)$ be the constant from Proposition 2.3, that is, every (c_1, c_2) -quasi-geodesic lies in a κ -neighborhood of a geodesic with the same endpoints and vice versa. Let $x_1 \in VG$ with $d(x_1, \pi_0) > 2\kappa$ and let $\beta_1 \in \Gamma$ with $x_0\beta_1 = x_1$. Then $\pi_1 := \pi_0\beta_1$ cannot lie 2κ -close to a geodesic between ω and η_0 . Thus, we have $\eta_1 := \eta_0\beta_1 \neq \eta_0$. Since α_0 is a hyperbolic element, so is $\alpha_1 := \beta_1^{-1}\alpha_0\beta_1$. Continuing like this we obtain the sequences as desired. Among the automorphisms α_i and α_i^{-1} we shall find a pair the product of which is non-trivial, elliptic, and fixes some vertex as required by the assertion.

Consider an infinite sequence $(B_i)_{i \in \mathbb{N}}$ of balls of radius 2κ around elements of π_0 that converge to ω . Since Γ acts transitively on VG , there is a finite number n such that each of these balls consists of n vertices. As the number of the bi-infinite quasi-geodesics with non-trivial intersection with B_i increases with i and tends to infinity, there is a ball B_m such that some vertex $b \in B_m$ lies on two distinct quasi-geodesics π_i, π_j with $i \neq j$. Since $d(x_k, x_k\alpha_k) = d(x_0, x_0\alpha_0)$ for all $k \in \mathbb{N}$ and since all balls of radius $d(x_0, x_0\alpha_0)$ have the same number of vertices, we may even assume that $b\alpha_i^{-1} = b\alpha_j^{-1}$. Let us consider the automorphism $\alpha_i^{-1}\alpha_j$. This automorphism obviously fixes b , so it is an elliptic element, and it is non-trivial because of $\alpha_i \neq \alpha_j$. This proves statement (ii).

It remains to prove (iii). We continue with the same notation as in the proof of (ii). Let $\gamma := \alpha_i^{-1}\alpha_j$ be the elliptic element we constructed in the proof of (ii). Then, for each $k \in \mathbb{N}$, $\gamma_k := \alpha^k\gamma\alpha^{-k}$ is an elliptic element that is not trivial but acts trivially on $b\alpha^{-k}$. By a similar argument as above, we shall find two automorphisms of the γ_k and γ_k^{-1} that will satisfy together with their product the assertion (iii). Each elliptic element γ_k has to act on the set of infinite (c_1, c_2) -quasi-geodesics from $b\alpha^{-k}$ to ω . Let us consider the sequence of balls $(D_k)_{k \in \mathbb{N}}$ with center $b\alpha^{-k}$ and radius 2κ . Like in the proof of (ii), there is an $m \in \mathbb{Z}$ such that two distinct

γ_k, γ_l , with $k \neq l$, both fix a vertex $y \in D_m$. Then $\gamma_k^{-1}\gamma_l$ also fixes y and it is again non-trivial because $\gamma_k \neq \gamma_l$. Hence $\gamma_k^{-1}\gamma_l$ satisfies assertion (iii). \square

With this information about hyperbolic and elliptic elements in automorphism groups of hyperbolic graphs we can now prove our main result.

Theorem 3.4. *For every planar one-ended hyperbolic graph G , and every group Γ of automorphisms of G that acts transitively on VG , no hyperbolic boundary point of G is fixed by all elements of Γ .*

Proof. Let us suppose, seeking for a contradiction, that there is a planar one-ended hyperbolic graph G and a subgroup Γ of $\text{Aut}(G)$ acting transitively on VG and fixing a hyperbolic boundary point ω . Let δ be the hyperbolicity constant of G as above, and let d be the degree of some, and hence any vertex of G . Then we have $d \geq 3$ and, by Lemma 2.6 and Theorem 2.5, every automorphism in Γ is consistent, either spin-preserving or spin-reversing. Let uvw be a 3-vertex subpath of a path P . We say that a vertex $x \in N(v) \setminus \{u, w\}$ lies *to the right* of P if in the spin of v we have vx between vw and vu . If x does not lie to the right of P then it lies *to the left* of P .

By Lemma 3.3 (iii) there are three non-trivial elliptic elements such that one of them is the product of the other two. If all non-trivial elliptic automorphisms are spin-reversing, then this is a direct contradiction as the product of any two spin-reversing automorphisms has to be spin-preserving. Thus, we may assume that an elliptic element $\varphi \in \Gamma$ is spin-preserving.

Let y be a vertex with $y\varphi \neq y$. As φ is elliptic, there is a minimal $n \in \mathbb{N}$ such that $y\varphi^n = y$. For all $i = 0, \dots, n-1$, let g_i be a geodesic from $y_i := y\varphi^i$ to y_{i+1} such that $g_i\varphi = g_{i+1}$ and let π_i be an infinite geodesic from $y\varphi^i$ converging to ω such that $\pi_i\varphi = \pi_{i+1}$. Then $C := g_0 \dots g_{n-1}$ is a cycle of length $n \cdot l(g_0)$. We distinguish two cases that will both lead to a contradiction: either π_i and π_{i+1} intersect infinitely often or not.

Let us first consider the case that they have only finitely many common vertices. By choosing some other vertex y' on π_0 instead of y we may assume that the corresponding geodesics $y_i\pi_i$ and $y_{i+1}\pi_{i+1}$ have no common vertex. In this situation, we shall show that any hyperbolic boundary point but ω is separated from ω by some finite cycle, which is impossible as G has precisely one end.

For every vertex on π_0 with distance larger than $l(g_0) + \delta$ to y_0 there is a vertex on π_1 of distance at most δ . We fix a geodesic between each such two vertices whose intersection with π_0, π_1 , respectively, is a connected subpath. If we consider an infinite sequence $(z_i)_{i \in \mathbb{N}}$ of vertices of π_0 with strictly increasing distance to y_0 , then either there are two infinite subsequences such that the chosen geodesics for one of them always lie to the right of π_0 for each vertex and for the other sequence always to the left of π_0 , or the geodesics for all but finitely many lie to the same side, say to the right of π_0 . If the first of these two situations occurs, then any infinite geodesic either is equivalent to π_0 —and hence converges to ω —or is separated by some finite cycle from ω , which is impossible as G is one-ended. Thus, we may assume that all the above described geodesics lie eventually to the right of π_0 . Let V_0 be a subset of VG consisting of all the vertices from the paths π_0, π_1, g_0 and from all the paths from π_0 whose first vertex lies to the right of π_0 and that has only vertices not in $\pi_0 \cup \pi_1 \cup g_0$ except for its first vertex. This means that V_0

consists of all those vertices of G that are separated in the plane by $\pi_0 \cup \pi_1 \cup g_0$ from any vertex that lies to the right of π_1 . Then any ray in $G[V_0]$ has to converge to ω . Similarly we find V_1, \dots, V_{n-1} , always taking the vertices to the right of π_i to obtain V_i , because φ is spin-preserving. We conclude that any other hyperbolic boundary point is separated from ω by C , which is impossible since G has precisely one end.

Thus, the only case left is that π_0 and π_1 have infinitely many common vertices. By Proposition 2.2, there is a $k \in \mathbb{N}$ such that for all but finitely many vertices x on π_0 we have $d(x, x\varphi) \leq k + 2\delta$. Again we distinguish two cases: either $\pi_0 - \pi_1$ contains infinitely many vertices or only finitely many. Let us first suppose that there are infinitely many vertices in $\pi_0 - \pi_1$. Let $(x_i)_{i \in \mathbb{N}}$ be a sequence of pairwise distinct vertices on $\pi_0 \cap \pi_1$ such that the predecessor of x_i on π_0 does not lie on π_1 and such that the predecessor of x_i on π_1 lies always to the same side of π_0 at the vertex x_i , say to the right. Then there is an $M \in \mathbb{N}$ such that we have for all $i \geq M$ that $C_i := x_i \pi_1 x_i \varphi \pi_2 \dots x_i \varphi^{n-1} \pi_0 x_i$ is a cycle separating C from ω . The inequality $d(x, x\varphi) \leq k + 2\delta$ immediately implies that all these cycles have length at most $2n(k + 2\delta)$ because of $d(x_i, x_i \varphi^{n-1}) \leq n(k + 2\delta)$. Now consider the ball B with center x_1 and radius $2n(k + 2\delta)$. As G is locally finite, $G - B$ has only finitely many components and precisely one of them is infinite because G is one-ended. Let N denote the number of vertices in finite components of $G - B$. Then we look at any ball B' with center x_i for an $i > N + 2n(k + 2\delta) + 1$ and radius $2n(k + 2\delta)$. Again, $G - B'$ has only one infinite component and the number of vertices in the finite components of $G - B'$ is precisely N by the transitivity of G . But because of $C_i \subseteq B'$, the component A that contains x_1 is finite. Since π_0 is a geodesic, A contains all x_j with $j \leq N + 1$, so there are at least $N + 1$ vertices in finite components of $G - B'$ which is a contradiction to the transitivity of Γ on G .

Thus, the only remaining case is when there are only finitely many vertices in $\pi_0 - \pi_1$. By replacing y by another suitable vertex on π_0 , we may assume that all the vertices of π_0 lie on π_1 or vice versa. But then either

$$\pi_n = y_0 \pi_n y_{n-1} \dots y_1 \pi_1 y_0 \pi_0$$

or

$$\pi_0 = y_0 \pi_0 y_1 \pi_1 \dots y_{n-1} \pi_{n-1} y_0 \pi_n$$

contains a cycle and so it cannot be an infinite geodesic, contrary to our assumption. This completes the proof. \square

REFERENCES

- [1] J. ALONSO ET AL., Notes on word hyperbolic groups, in: E. GHYS, A. HAEFLIGER, AND A. VERJOVSKY (eds.), *Group Theory from a Geometrical Viewpoint* (Trieste, 1990), pp. 3–63, World Scientific, 1991.
- [2] L. BABAI, The growth rate of vertex-transitive planar graphs, in: *Proceedings of the Eighth Annual ACM-SIAM Symposium on Discrete Algorithms* (New Orleans, LA, 1997), pp. 564–573, ACM, New York, 1997.
- [3] L. BABAI AND M. WATKINS, Connectivity of infinite graphs having a transitive torsion group action, *Arch. Math. (Basel)*, **34**, 90–96 (1980).
- [4] M. COORNAERT, T. DELZANT, AND A. PAPADOPOULOS, Notes sur les groupes hyperboliques de Gromov, *Lecture Notes in Mathematics*, volume 1441, Springer-Verlag, 1990.
- [5] R. DIESTEL, *Graph Theory (4th edition)*, Springer-Verlag, 2010, Electronic edition available at: <http://www.math.uni-hamburg.de/home/diestel/books/graph.theory>.

- [6] R. DIESTEL, H. JUNG, AND R. MÖLLER, On vertex transitive graphs of infinite degree, *Arch. Math. (Basel)*, **60**, 591–600 (1993).
- [7] E. GHYS AND P. DE LA HARPE, Sur les groupes hyperboliques, d’après M. Gromov, *Progress in Math.*, volume 83, Birkhäuser, Boston, 1990.
- [8] M. GROMOV, Hyperbolic Groups, in: S. GERSTEN (ed.), *Essays in group theory, MSRI*, volume 8, pp. 75–263, Springer, New York, 1987.
- [9] W. IMRICH, On Whitney’s theorem on the unique embeddability of 3-connected planar graphs., in: *Recent Adv. Graph Theory, Proc. Symp. Prague 1974*, pp. 303–306, 1975.
- [10] V. KAIMANOVICH AND W. WOESS, Boundary and entropy of space homogeneous Markov chains, *Ann. Probab.*, **30**, 323–363 (2002).
- [11] R. MÖLLER, Ends of graphs II, *Math. Proc. Cambridge Phil. Soc.*, **111**, 455–460 (1992).
- [12] H. WHITNEY, Congruent Graphs and the Connectivity of Graphs, *American J. of Mathematics*, **54**, 150–168 (1932).
- [13] W. WOESS, Topological groups and infinite graphs, *Discrete Math.*, **95**, 373–384 (1991).
- [14] W. WOESS, Fixed sets and free subgroups of groups acting on metric spaces, *Math. Z.*, **214**, 425–440 (1993).
- [15] W. WOESS, *Random walks on infinite graphs and groups*, Cambridge University Press, 2000.

TECHNISCHE UNIVERSITÄT GRAZ, STEYRERGASSE 30, 8010 GRAZ, AUSTRIA

Email address: `georgakopoulos@tugraz.at`

FACHBEREICH MATHEMATIK, UNIVERSITÄT HAMBURG, BUNDESSTRASSE 55, 20146 HAMBURG, GERMANY

Email address: `matthias.hamann@math.uni-hamburg.de`