

Forcing and Independence Proofs: Assignment 3

Part A: The forcing relation

1. Prove the following facts about the “semantic” forcing relation.¹
 - (a) Suppose that $\forall p \in \mathbb{P} \exists q \leq p (q \Vdash \phi)$. Then $\mathbf{1} \Vdash \phi$.
 - (b) Similarly, suppose $p_0 \in \mathbb{P}$ is such that $\forall p \leq p_0 \exists q \leq p (q \Vdash \phi)$. Then $p_0 \Vdash \phi$.
 - (c) The following are equivalent:
 - i. $p_0 \Vdash \phi$,
 - ii. $\forall p \leq p_0 (p \Vdash \phi)$, and
 - iii. $\{p \leq p_0 : p \Vdash \phi\}$ is dense below p_0 .
2. For $p \in \mathbb{P}$ and ϕ in the forcing language, we say p *decides* ϕ if $p \Vdash \phi$ or $p \Vdash \neg\phi$. Show that for every $p \in \mathbb{P}$ and every ϕ , there is $q \leq p$ which decides ϕ .
3. Prove the inductive rule for forcing a disjunction: $p \Vdash \phi \vee \psi$ iff $\{q \leq p : q \Vdash \phi \vee q \Vdash \psi\}$ is dense below p . You may do this either by reducing $\phi \vee \psi$ to a statement with conjunctions and negations, or use the definition of the semantic relation and prove the statement directly.

¹I.e., using the definition: $(p \Vdash \varphi)^M \Leftrightarrow \forall G \mathbb{P}$ -generic over M if $p \in G$ then $M[G] \models \varphi$, not using the “syntactic” forcing relation \Vdash^* .

Part B: Forcing

1. Let τ be a name such that $p \Vdash \tau \in \check{\omega}$. Show that there exists $q \leq p$ and $n \in \omega$ such that $q \Vdash \tau = \check{n}$. We say that q *decides* τ .
2. Consider the partial order $Fn(\omega, \omega)$, i.e., finite functions p with $\text{dom}(p), \text{ran}(p) \subseteq \omega$ ordered by $q \leq p$ iff $q \supseteq p$ (the standard partial order for adding a new real). This forcing is typically called *Cohen forcing*.

Let G be Cohen-forcing-generic over M . Show that $f_G := \bigcup G$ has the following property: for every $x \in \omega^\omega \cap M$, there are infinitely many $n \in \omega$, such that $x(n) < f_G(n)$ (we say that f_G is an *unbounded real over M*).

Hint: for every $x \in \omega^\omega \cap M$ and every $k \in \omega$, define appropriate dense sets $D_{x,k} = \dots$