

# AN EXCLUDED MINORS METHOD FOR INFINITE MATROIDS

Nathan Bowler      Johannes Carmesin

December 3, 2012

## Abstract

The notion of thin sums matroids was invented to extend the notion of representability to non-finitary matroids. A matroid is tame if every circuit-cocircuit intersection is finite. We prove that a tame matroid is a thin sums matroid over a finite field  $k$  if and only if all its finite minors are representable over  $k$ .

We expect that the method we use to prove this will make it possible to lift many theorems about finite matroids representable over a finite field to theorems about tame thin sums matroids over these fields. We give three examples of this: various characterisations of binary tame matroids and of regular tame matroids, and unique representability of ternary tame matroids.

## 1 Introduction

Given a family of vectors in a vector space over some field  $k$ , we get a matroid structure on that family whose independent sets are given by the linearly independent subsets of the family. Matroids arising in this way are called *representable* matroids over  $k$ . A classical theorem of Tutte [12] states that a finite matroid is binary (that is, representable over  $\mathbb{F}_2$ ) if and only if it does not have  $U_{2,4}$  as a minor. In the same spirit, a key aim of finite matroid theory has been to determine such ‘forbidden minor’ characterisations for the classes of matroids representable over other finite fields. For example Bixby and Seymour [2, 11] characterized the finite ternary matroids (those representable over  $\mathbb{F}_3$ ) by forbidden minors, and more recently there is a forbidden minors characterisation for the finite matroids representable over  $\mathbb{F}_4$ , due to Geelen, Gerards and Kapoor [8]. This remains an open problem for all other finite fields. In this paper, we consider the corresponding problem for infinite matroids.

It is clear that any representable matroid is *finitary*, that is, all its circuits are finite, and so many interesting examples of infinite matroids are not representable. However, since the construction of many standard examples, including the algebraic cycle matroids of infinite graphs, is suggestively similar to that of representable matroids, the notion of *thin sums matroids* was introduced in [4]:

it is a generalisation of representability which captures these infinite examples. We will work with thin sums matroids rather than with representable matroids.

In [1] it was shown that the class of tame thin sums matroids over a fixed field is closed under duality, where a matroid is *tame* if any circuit-cocircuit intersection is finite. On the other hand, there are thin sums matroids whose dual is not a thin sums matroid [3] - such counterexamples cannot be tame. A simple consequence of this closure under duality is that the class of tame thin sums matroids over a fixed field is closed under taking minors, and so we may consider the forbidden minors for this class.

Minor closed classes may have infinite ‘minimal’ forbidden minors. For example the class of finitary matroids has the infinite circuit  $U_{1,\mathbb{N}}^*$  as a forbidden minor. Similarly, the class of tame thin sums matroids over  $\mathbb{R}$  has  $U_{2,\mathcal{P}(\mathbb{R})}$  as a forbidden minor. However, our main result states that the class of tame thin sums matroids over a fixed *finite* field has only finite minimal forbidden minors.

**Theorem 1.1.** *Let  $M$  be a tame matroid and  $k$  be a finite field. Then  $M$  is a thin sums matroid over  $k$  if and only if every finite minor of  $M$  is  $k$ -representable.*

The proof is by a compactness argument. All previous compactness proofs in infinite matroid theory known to the authors use only that either all finite restrictions or all finite contractions have a certain property to conclude that the matroid itself has the desired property. For our purposes, arguments of this kind must fail because there is a tame matroid all of whose finite restrictions and finite contractions are binary but which is not a thin sums matroid over  $\mathbb{F}_2$  - in fact, it has a  $U_{2,4}$ -minor. We shall briefly sketch how to construct such a matroid. Start with  $U_{2,4}$ , and add infinitely many edges parallel to one of its edges. This ensures that every finite contraction is binary. If we also add infinitely many edges which are parallel in the dual to some other edge then we guarantee in addition that all finite restrictions are binary, but the matroid itself has a  $U_{2,4}$  minor.

Theorem 1.1 implies that each of the excluded minor characterisations for finite representable matroids mentioned in the first paragraph extends to tame matroids. Thus, for example, a tame matroid is a thin sums matroid over  $\mathbb{F}_2$  if and only if it has no  $U_{2,4}$  minor. Any future excluded minor characterisations for finite matroids representable over a fixed finite field will also immediately extend to tame matroids by this theorem.

We expect that our approach will make it possible to lift many other standard theorems about finite matroids representable over a finite field to theorems about tame thin sums matroids over the same field. We shall give four instances of this. Firstly, as for finite binary matroids, we obtain nine equivalent simple characterizations of tame thin sums matroids over  $\mathbb{F}_2$ . Secondly, we show that a tame matroid is regular (that is, a thin sums matroid over every field) if and only if all its finite minors are, and that regularity is equivalent to signability for tame matroids (see Section 5 for a definition). Thirdly, we introduce an extension of the notion of uniqueness of representations for finite matroids to the infinite setting, and prove that tame thin sums matroids over  $\mathbb{F}_3$  have unique

representations in this sense. A consequence of this is that the finite, algebraic and topological cycle matroids of a given graph each have a unique signing. We illustrate how these signings encode some of the structure of the graph. Fourthly, we show that the characterisations of representability over various sets of finite fields in terms of representability over partial fields extend in a uniform way to infinite tame matroids.

Our method applies to excluded minor characterisations of properties other than representability. In [9] the same method is employed to show that the tame matroids all of whose finite minors are graphic are precisely those matroids that arise from some *graph-like space*, in the sense that the circuits are given by topological circles and the cocircuits by topological bonds.

The paper is organized as follows. After the preliminaries in Section 2, in Section 3 we examine in detail the special case of binary matroids. Although this case is simpler than the general case, it illustrates some ideas that will be important in the proof of the main result. At the end of Section 3, we offer some related open problems. In Section 4 we prove our main result. In Section 5 we discuss the implications for regular and ternary matroids, and matroids representable over partial fields.

## 2 Preliminaries

### 2.1 Basics

Throughout, notation and terminology for graphs are those of [7], and for matroids those of [10, 5].  $M$  always denotes a matroid and  $E(M)$  (or just  $E$ ),  $\mathcal{I}(M)$  and  $\mathcal{C}(M)$  denote its ground set and its sets of independent sets and circuits, respectively.

A set system  $\mathcal{I} \subseteq \mathcal{P}(E)$  is the set of independent sets of a matroid iff it satisfies the following *independence axioms* [5].

- (I1)  $\emptyset \in \mathcal{I}(M)$ .
- (I2)  $\mathcal{I}(M)$  is closed under taking subsets.
- (I3) Whenever  $I, I' \in \mathcal{I}(M)$  with  $I'$  maximal and  $I$  not maximal, there exists an  $x \in I' \setminus I$  such that  $I + x \in \mathcal{I}(M)$ .
- (IM) Whenever  $I \subseteq X \subseteq E$  and  $I \in \mathcal{I}(M)$ , the set  $\{I' \in \mathcal{I}(M) \mid I \subseteq I' \subseteq X\}$  has a maximal element.

A set system  $\mathcal{C} \subseteq \mathcal{P}(E)$  is the set of circuits of a matroid iff it satisfies the following *circuit axioms* [5].

- (C1)  $\emptyset \notin \mathcal{C}$ .
- (C2) No element of  $\mathcal{C}$  is a subset of another.

- (C3) (Circuit elimination) Whenever  $X \subseteq o \in \mathcal{C}(M)$  and  $\{o_x \mid x \in X\} \subseteq \mathcal{C}(M)$  satisfies  $x \in o_y \Leftrightarrow x = y$  for all  $x, y \in X$ , then for every  $z \in o \setminus (\bigcup_{x \in X} o_x)$  there exists a  $o' \in \mathcal{C}(M)$  such that  $z \in o' \subseteq (o \cup \bigcup_{x \in X} o_x) \setminus X$ .
- (CM)  $\mathcal{I}$  satisfies (IM), where  $\mathcal{I}$  is the set of those subsets of  $E$  not including an element of  $\mathcal{C}$ .

**Lemma 2.1.** *Let  $M$  be a matroid and  $s$  be a base. Let  $o_e$  and  $b_f$  a fundamental circuit and a fundamental cocircuit with respect to  $s$ , then*

1.  $o_e \cap b_f$  is empty or  $o_e \cap b_f = \{e, f\}$  and
2.  $f \in o_e$  iff  $e \in b_f$ .

*Proof.* To see the first note that  $o_e \subseteq s + e$  and  $b_f \subseteq (E \setminus s) + f$ . So  $o_e \cap b_f \subseteq \{e, f\}$ . As a circuit and a cocircuit can never meet in only one edge, the assertion follows.

To see the second, first let  $f \in o_e$ . Then  $f \in o_e \cap b_f$ , so by (1)  $o_e \cap b_f = \{e, f\}$  and so  $e \in b_f$ . The converse implication is the dual statement of the above implication.  $\square$

**Lemma 2.2.** *For any circuit  $o$  containing two edges  $e$  and  $f$ , there is a cocircuit  $b$  such that  $o \cap b = \{e, f\}$ .*

*Proof.* As  $o - e$  is independent, there is a base including  $o - e$ . By Lemma 2.1, the fundamental cocircuit of  $f$  of this base intersects  $o$  in  $e$  and  $f$ , as desired.  $\square$

**Lemma 2.3.** *Let  $M$  be a matroid with ground set  $E = C \dot{\cup} X \dot{\cup} D$  and let  $o'$  be a circuit of  $M' = M/C \setminus D$ . Then there is an  $M$ -circuit  $o$  with  $o' \subseteq o \subseteq o' \cup C$ .*

*Proof.* Let  $s$  be any  $M$ -base of  $C$ . Then  $s \cup o'$  is  $M$ -dependent since  $o'$  is  $M'$ -dependent. On the other hand,  $s \cup o' - e$  is  $M$ -independent whenever  $e \in o'$  since  $o' - e$  is  $M'$ -independent. Putting this together yields that  $s \cup o'$  contains an  $M$ -circuit  $o$ , and this circuit must not avoid any  $e \in o'$ , as desired.  $\square$

**Corollary 2.4.** *Let  $M'$  be a minor of  $M$ . Further let  $o'$  be an  $M'$ -circuit and  $b'$  be an  $M'$ -cocircuit. Then there is an  $M$ -circuit  $o \subseteq o' \cup (E(M) \setminus E(M'))$  and an  $M$ -cocircuit  $b \subseteq b' \cup (E(M) \setminus E(M'))$  such that  $o \cap b = o' \cap b'$ .*

*Proof.* Extend  $o'$  using contracted edges only and  $b'$  using deleted edges only as in Lemma 2.3 and its dual.  $\square$

## 2.2 Scrawls

A *scrawl* is a union of circuits. For any matroid  $M$ ,  $M$  can be recovered from its set of scrawls since the circuits are precisely the minimal nonempty scrawls.

We can formulate the matroid axioms in terms of scrawls. The scrawl axioms for a set  $\mathcal{S} \subseteq \mathcal{P}(E)$  are the following.

- (S1) Any union of elements of  $\mathcal{S}$  is in  $\mathcal{S}$ .

- (S2) (scrawl elimination) Whenever  $X \subseteq w \in \mathcal{S}$  and  $\{w_x \mid x \in X\} \subseteq \mathcal{S}$  satisfies  $x \in w_y \Leftrightarrow x = y$  for all  $x, y \in X$ , then for every  $z \in w \setminus (\bigcup_{x \in X} w_x)$  there exists a  $w' \in \mathcal{S}$  such that  $z \in w' \subseteq (w \cup \bigcup_{x \in X} w_x) \setminus X$ .
- (SM)  $\mathcal{I}$  satisfies (IM), where  $\mathcal{I}$  is the set of those subsets of  $E$  not including a nonempty element of  $\mathcal{S}$ .

**Proposition 2.5.** *Let  $\mathcal{S} \subseteq \mathcal{P}(E)$ .  $\mathcal{S}$  is the set of scrawls of a matroid with ground set  $E$  if and only if it satisfies the scrawl axioms.*

*Proof.* Clearly, (S1) and (SM) are satisfied by the set of scrawls of any matroid  $M$ . Now let  $w, X, \{w_x \mid x \in X\}$  and  $z$  as in (S2). Let  $o$  be an  $M$ -circuit such that  $z \in o \subseteq w$ , which exists as  $w$  is a union of circuits. Let  $\bar{X} = o \cap X$ . For every  $x \in \bar{X}$ , let  $o_x$  be an  $M$ -circuit such that  $x \in o_x \subseteq w_x$ . Now apply circuit elimination to obtain a circuit  $o'$  such that  $z \in o' \subseteq (o \cup \bigcup_{x \in \bar{X}} w_x) \setminus \bar{X}$ . By the choice of the  $w_x$  every  $x \in X \setminus \bar{X}$  is not in  $o \cup \bigcup_{x \in \bar{X}} w_x$ . This completes the proof of (S2).

Conversely, let  $\mathcal{S}$  satisfy the scrawl axioms and let  $\mathcal{C}$  be the set of minimal nonempty elements of  $\mathcal{S}$ . First we show that any nonempty element of  $\mathcal{S}$  is a union of elements of  $\mathcal{C}$ . Let  $w \in \mathcal{S}$  and  $z \in w$  be given. It suffices to show that there is a minimal nonempty element  $o$  of  $\mathcal{S}$  such that  $z \in o \subseteq w$ . By (SM) there is a maximal  $I \in \mathcal{I}$  such that  $I \subseteq w - z$ . Let  $X = (w - z) \setminus I$ . For every  $x \in X$ , there is by maximality of  $I$  an element  $w_x$  of  $\mathcal{S}$  included in  $I + x$ . Now apply scrawl elimination to obtain an element  $w'$  of  $\mathcal{S}$  with  $z \in w' \subseteq I + z$ .

We show  $w' \in \mathcal{C}$ . Suppose for a contradiction that  $w'$  properly includes a nonempty element  $w''$  of  $\mathcal{S}$ . Note that  $z \in w''$  because otherwise  $I$  would include a nonempty element of  $\mathcal{S}$ , which is impossible. Pick  $x \in w' \setminus w''$ . Now we may apply scrawl elimination (with  $x$  in place of  $z$  and  $\{z\}$  in place of  $X$ ) to obtain an element of  $\mathcal{S}$  included in  $I$  which is nonempty as it contains  $x$ . This is a contradiction.

Second, we show that  $\mathcal{C}$  satisfies the circuit axioms. (C1),(C2) and (CM) are clear. Now let  $o, X, \{o_x \mid x \in X\}$  and  $z$  as in (C3). We apply (S3) and obtain a  $w' \in \mathcal{S}$  such that  $z \in w' \subseteq (o \cup \bigcup_{x \in X} o_x) \setminus X$ . Above we showed how we find an  $o' \in \mathcal{C}$  such that  $z \in o' \subseteq w'$ , proving (C3).

Now let  $M$  be the matroid whose circuits are given by  $\mathcal{C}$ . Then the set of scrawls of  $M$  is  $\mathcal{S}$  by (S1).  $\square$

Dually a *coscrawl* is a union of cocircuits. Since no circuit and cocircuit can meet in only one element, no scrawl and coscrawl can meet in only one element. In fact, this property gives us a simple characterisation of scrawls in terms of coscrawls and vice versa.

**Lemma 2.6.** *Let  $M$  be a matroid, and let  $w \subseteq E$ . The following are equivalent:*

1.  $w$  is a scrawl of  $M$ .
2.  $w$  never meets a cocircuit of  $M$  just once.
3.  $w$  never meets a coscrawl of  $M$  just once.

*Proof.* It is clear that (1) implies (3) and (3) implies (2), so it suffices to show that (2) implies (1). Suppose that (2) holds and let  $e \in w$ . Then in the minor  $M/(w-e) \setminus (E \setminus w)$  on the groundset  $\{e\}$ ,  $e$  cannot be a co-loop, by the dual of Lemma 2.3 and (2). So  $e$  must be a loop, and by Lemma 2.3 there is a circuit  $o_e$  with  $e \in o_e \subseteq w$ . Thus  $w$  is the union of the  $o_e$ , and so is a scrawl.  $\square$

**Corollary 2.7.** *Let  $M$  be a matroid with ground set  $E = C \dot{\cup} X \dot{\cup} D$ , and let  $w' \subseteq X$ . Then  $w'$  is a scrawl of  $M' = M/C \setminus D$  if and only if there is a scrawl  $w$  of  $M$  with  $w' \subseteq w \subseteq w' \cup C$ .*

*Proof.* The ‘only if’ direction is clear from Lemma 2.3 and the definition of a scrawl. Conversely, if there is such a  $w$  then by the dual of Lemma 2.3  $w'$  can never meet a cocircuit of  $M'$  just once, so by Lemma 2.6 it is a scrawl of  $M'$ .  $\square$

## 2.3 Thin sums matroids

Throughout the whole paper, we will follow the convention that if we write that a sum equals zero then this implicitly includes the statement that this sum is well-defined, that is, that only finitely many summands are nonzero.

**Definition 2.8.** Let  $A$  be a set, and  $k$  a field. Let  $f = (f_e | e \in E)$  be a family of functions from  $A$  to  $k$ , and let  $\lambda = (\lambda_e | e \in E)$  be a family of elements of  $k$ . We say that  $\lambda$  is a *thin dependence* of  $f$  iff for each  $a \in A$  we have

$$\sum_{e \in E} \lambda_e f_e(a) = 0,$$

We say that a subset  $I$  of  $E$  is *thinly independent* for  $f$  iff the only thin dependence of  $f$  which is 0 everywhere outside  $I$  is  $(0 | e \in E)$ . The *thin sums system*  $M_f$  of  $f$  is the set of such thinly independent sets. This isn’t always the set of independent sets of a matroid [5], but when it is we call it the *thin sums matroid* of  $f$ .

This definition is deceptively similar to the definition of the representable matroid corresponding to  $f$  considered as a family of vectors in the  $k$ -vector space  $k^A$ . The difference is in the more liberal definition of dependence: it is possible for  $\lambda$  to be a thin dependence even if there are infinitely many  $e \in E$  with  $\lambda_e \neq 0$ , provided that for each  $a \in A$  there are only finitely many  $e \in E$  such that *both*  $\lambda_e \neq 0$  and  $f_e(a) \neq 0$ .

Indeed, the notion of thin sums matroid was introduced as a generalisation of the notion of representable matroid: every representable matroid is finitary, but this restriction does not apply to thin sums matroids.

There are many natural examples of thin sums matroids: for example, the algebraic cycle matroid of any graph not including a subdivision of the Bean graph is a thin sums matroid, as follows:

**Definition 2.9.** Let  $G$  be a graph with vertex set  $V$  and edge set  $E$ , and  $k$  a field. We can pick a direction for each edge  $e$ , calling one of its ends its *source*  $s(e)$  and the other its *target*  $t(e)$ . Then the family  $f^G = (f_e^G | e \in E)$  of functions from  $V$  to  $k$  is given by  $f_e = \chi_{t(e)} - \chi_{s(e)}$ , where for any vertex  $v$  the function  $\chi_v$  takes the value 1 at  $v$  and 0 elsewhere.

**Theorem 2.10** ([5, §6.7]). *Let  $G$  be a graph not including any subdivision of the Bean graph. Then  $M_{f^G}$  is the algebraic cycle matroid of  $G$ .*

The following connection between scrawls and linear dependences will turn out to be useful.

**Lemma 2.11.** *Let  $M(f)$  be a finite representable matroid, and let  $c$  be a linear dependence for  $f$ . Then the support of  $c$  is a scrawl.*

*Proof.* If  $c$  is a linear dependence, then its support is either empty or includes a circuit  $o$ . Then there is a linear dependence  $c_o$  with support precisely  $o$ . Pick  $e \in o$ , then  $c - c_o \cdot \frac{c(e)}{c_o(e)}$  is also a linear dependence whose support is a proper subset of the support of  $c$ . If we repeat this process once for every element in the support of  $c$ , then we end up with the empty set and have constructed for every element in the support of  $c$  a circuit that contains this element and that also is contained in the support of  $c$ .  $\square$


This Lemma is also true for infinite tame thin sums matroids [1].

### 3 Binary matroids

**Theorem 3.1.** *Let  $M$  be a tame matroid. Then the following are equivalent:*

1.  $M$  is a binary thin sums matroid.
2. For any circuit  $o$  and cocircuit  $b$  of  $M$ ,  $|o \cap b|$  is even.
3. For any circuit  $o$  and cocircuit  $b$  of  $M$ ,  $|o \cap b| \neq 3$
4.  $M$  has no minor isomorphic to  $U_{2,4}$ .
5. If  $o_1, o_2$  are circuits then  $o_1 \Delta o_2$  is empty or includes a circuit.
6. If  $o_1, o_2$  are circuits then  $o_1 \Delta o_2$  is a disjoint union of circuits.
7. If  $(o_i | i \in I)$  is a finite family of circuits then  $\Delta_{i \in I} o_i$  is empty or includes a circuit.
8. If  $(o_i | i \in I)$  is a finite family of circuits then  $\Delta_{i \in I} o_i$  is a disjoint union of circuits.
9. For any base  $s$  of  $M$ , and any circuit  $o$  of  $M$ ,  $o = \Delta_{e \in o \setminus s} o_e$ , where  $o_e$  is the fundamental circuit of  $e$  with respect to  $s$ .

*Proof.* We shall prove the following implications:



Those implications indicated by dotted arrows are clear. We shall prove the remaining implications.

(2) implies (1): We need to find a suitable thin sums system. Let  $A$  be the set of cocircuits of  $M$ , and let  $E \xrightarrow{f} \mathbb{F}_2^A$  be the map sending  $e$  to the function which sends  $b \in A$  to 1 if  $e \in b$  and 0 otherwise.

We are to show that the thin sums matroid  $M_{ts}$  defined by  $f$  is  $M$ . Since the characteristic function of any  $M$ -circuit is a thin dependence for  $f$  with support equal to that circuit by (2), any  $M$ -dependent set is also  $M_{ts}(f)$ -dependent.

It remains to show that the support of every non-zero thin dependence is  $M$ -dependent. By Lemma 2.6 the support of every non-zero thin dependence includes a circuit, as desired.

(2) implies (8): Let  $(o_i | i \in I)$  be a finite family of circuits. By Zorn's Lemma, we can choose a maximal family  $(o_j | j \in J)$  of disjoint circuits such that  $\bigcup_{j \in J} o_j \subseteq \Delta_{i \in I} o_i$ , and let  $w = \Delta_{i \in I} o_i \setminus \bigcup_{j \in J} o_j$ . Let  $b$  be any cocircuit of  $M$ , so that  $|b \cap o_i|$  is even for each  $i \in I$ . Then  $|b \cap \Delta_{i \in I} o_i|$  is also even, and in particular finite. Since the  $o_j$  are disjoint, there can only be finitely many of them that meet  $b \cap \Delta_{i \in I} o_i$ , and since for each such  $j$  we have that  $|b \cap o_j|$  is even, it follows that  $|b \cap w|$  is even. In particular,  $b \cap w$  doesn't have just one element. Since  $b$  was arbitrary, by Lemma 2.6  $w$  is a scrawl of  $M$  and so if it is nonempty it includes a circuit. But in that case, we could add that circuit to the family  $(o_j | j \in J)$ , contradicting the maximality of that family. Thus  $w$  is empty, and  $\Delta_{i \in I} o_i = \bigcup_{j \in J} o_j$  is a disjoint union of circuits.

(5) implies (3): Suppose, for a contradiction, that (5) holds but (3) fails, and choose a circuit  $o$  and a cocircuit  $b$  with  $o \cap b = \{x, y, z\}$  of size 3. Pick a base  $s$  of  $(E \setminus b) + x$  including  $o - y - z$ , which exists by  $(IM)$ . As  $b$  is a cocircuit,  $b - x$  avoids some  $M$ -base, thus  $(E \setminus b) + x$  is spanning and thus  $s$  is spanning, as well. Let  $o_y$  and  $o_z$  be the fundamental circuits of  $y$  and  $z$  with respect to  $s$ .

It suffices to show that  $o_y \Delta o_z \subseteq o - x$ . Indeed, since  $y, z \in o_y \Delta o_z$ , (5) then yields a circuit properly included in  $o$ , which is impossible. By Lemma 2.2 we can't have  $o_y \cap b = \{y\}$  so we must have  $x \in o_y$ . Similarly,  $x \in o_z$ , and so  $x \notin o_y \Delta o_z$ . So it is sufficient to show that  $o_y$  and  $o_z$  agree outside  $o$ , in other words:  $o_y \subseteq o_z \cup o$  and  $o_z \subseteq o_y \cup o$ .



To see this, first note that by uniqueness of the fundamental circuit of  $y$  it suffices to show that  $y$  is spanned by  $(o_z - z) \cup (o - y - z)$ . As  $z$  is spanned by  $(o_z - z)$ ,  $o - y$  is spanned by  $(o_z - z) \cup (o - y - z)$ . Since  $o$  is a circuit,  $y$  is also spanned by  $(o_z - z) \cup (o - y - z)$ , as desired. A similar argument yields  $o_z \subseteq o_y \cup o$ , completing the proof.

(3) implies (4): Since any subset of the ground set of  $U_{2,4}$  of size 3 is both a circuit and a cocircuit, it is easy to find a circuit and cocircuit in  $U_{2,4}$  whose intersection has size 3. So we simply apply Corollary 2.4.

(4) implies (2): Suppose for a contradiction that (4) holds but (2) does not. Then let  $o$  be a circuit and  $b$  a cocircuit such that  $|o \cap b| = k$  is odd. By contracting  $o \setminus b$  and deleting  $b \setminus o$ , we obtain a minor  $M'$  of  $M$  in which  $o \cap b$  is both a circuit and a cocircuit. Let  $s$  be a minimal spanning set containing  $o \cap b$ , which exists by  $(IM^*)$ . Then in the minor  $M''$  of  $M'$  obtained by contracting  $s \setminus (o \cap b)$ ,  $(o \cap b)$  is spanning, and is still both a circuit and a cocircuit. By a similar removal, we can find a minor  $M'''$  of  $M''$  in which  $o \cap b$  is a circuit and a cocircuit and is both spanning and cospanning. Let  $x \in o \cap b$ . Then  $o \cap b - x$  is both a base and a cobase of  $M'''$ , and it is finite (it has size  $k - 1$ ). As  $o \cap b - x$  is a base and a cobase, the complement of  $o \cap b - x$  is also a base and a cobase. Thus the ground set of  $M'''$  is also finite (it has size  $2k - 2$ ). Applying the finite version of the theorem, then,  $M'''$  contains a  $U_{2,4}$  minor, which is also a minor of  $M$ , giving the desired contradiction.

(9) implies (2): first we will show that the following implies (2):

For any base  $s$  of  $M$ , any circuit  $o$  meets every fundamental cocircuit of  $s$  in an even number of edges. ( $\diamond$ )

To see that ( $\diamond$ ) implies (2), it suffices to show that every cocircuit  $b$  is fundamental cocircuit of some base  $s$ . Let  $e \in b$ . Then as  $b$  is a cocircuit,  $E \setminus (b - e)$  is spanning. Thus by  $(IM)$  there is a base  $s$  of  $E \setminus (b - e)$ , which clearly has  $b$  as fundamental cocircuit.

So it remains to see that (9) implies ( $\diamond$ ). By (9),  $o = \Delta_{e \in o \setminus s} o_e$ . Let  $b_f$  be some fundamental cocircuit of  $s$  for some  $f \in s$ . By Lemma 2.1  $o_e \cap b_f$  is empty or  $o_e \cap b_f = \{e, f\}$ . So it suffices to show that every  $f$  is in only finitely many  $o_e$ , which follows from the fact that  $o = \Delta_{e \in o \setminus s} o_e$  is well defined at  $f$ . This completes the proof.

(2) implies (9): we have to show for every edge  $f$  that it is contained in only finitely many  $o_e$  and that  $f \in o \iff f \in \Delta_{e \in o \setminus s} o_e(f)$ . If  $f \notin s$ , this is easy, so let  $f \in s$ . By Lemma 2.1  $f \in o_e$  iff  $e \in b_f$ . As  $M$  is tame  $|o \cap b_f|$  is finite, so there are only finitely many such  $e$ . By (2),  $|o \cap b_f|$  is even. If  $f \notin o$ , all such  $e$  are not contained in  $s$ , so  $f \notin \Delta_{e \in o \setminus s} o_e$ . If  $f \in o$ , all such  $e$  but  $f$  are not contained in  $s$ , so  $f \in \Delta_{e \in o \setminus s} o_e$ . This completes the proof.  $\square$

We remark that we might also put the duals of the statements in the list onto the list. It might be worth noting that (7) becomes false if we also allow  $I$  to be infinite. To see this, consider the finite cycle matroid of the graph obtained from a ray by adding a vertex that is adjacent to every vertex on the ray. Indeed, the symmetric difference of all 3-cycles is a ray starting at this new vertex. This

set is not empty, and nor does it include a circuit, so the infinite version of (7) fails.

More generally, the finite cycles of a locally finite graph generate the cycle space, which may contain infinite cycles [7].

We offer the following related open questions. Let (10) be the statement like (9) but for only one base of  $M$ . For finite matroids, (10) is equivalent to (9). Is the same true for tame matroids?

The following simple question also remains open:

**Open Question 3.2.** *Is every binary thin sums matroid tame?*

In a binary tame matroid, it is easy to see that any set meeting every cocircuit not in an odd number of edges is a disjoint union of circuits provided that the set is either countable or does not meet any cocircuit infinitely. A well-known result of Nash-Williams says that the above is also true if the matroid is the finite cycle matroid of some graph. Does this extend to all binary tame matroids?

**Open Question 3.3.** *Let  $M$  be a binary tame matroid and let  $X$  be a set that meets no cocircuit in an odd number of edges. Must  $X$  be a disjoint union of circuits?*

## 4 Representable matroids

The aim of this section is to provide an excluded-minors characterisation of thin sums matroids in the class of tame matroids. The following definition will be essential.

Let  $k$  be a field and let  $k^*$  denote the set of nonzero elements of  $k$ . A  $k$ -painting for the matroid  $M$  is a choice of functions  $c_o: o \rightarrow k^*$  for each circuit  $o$  of  $M$  and for each cocircuit  $b$  of  $M$  a function  $d_b: b \rightarrow k^*$  such that for any circuit  $o$  and cocircuit  $b$  we have

$$\sum_{e \in o \cap b} c_o(e) d_b(e) = 0. \tag{1}$$

A matroid is  $k$ -paintable if it has a  $k$ -painting. The method we will use is motivated by the following result, which is taken from [1]:

**Lemma 4.1.** *Let  $M$  be a tame matroid. Then  $M$  is a thin sums matroid over the field  $k$  iff  $M$  is  $k$ -paintable.*

Proving this Lemma would go beyond the scope of this paper, but we shall not need its full power here. Instead, we shall rely on the following weaker statement:

**Lemma 4.2.** *Let  $M$  be a finite matroid which is representable over the field  $k$ . Then  $M$  is  $k$ -paintable.*

*Proof.* Let  $M$  be given as  $M(\phi)$  for some  $E \xrightarrow{\phi} V$ . For each circuit  $o$  of  $M$ , pick some linear dependence  $\hat{c}_o$  of  $\phi$  with support  $o$ , and let  $c_o = \hat{c}_o \upharpoonright_o$ . Now let  $b$  be any cocircuit of  $M$ , and fix some  $e_b \in b$ . By Lemma 2.2, we can find for each  $e \in b - e_b$  some circuit  $o(e)$  of  $M$  such that  $o(e) \cap b = \{e_b, e\}$ . We define the map  $d_b: b \rightarrow k^*$  to be 1 at  $e_b$  and  $-\frac{c_o(e)(e_b)}{c_o(e)(e)}$  for  $e \in b - e_b$  (note that this choice ensures that (2) holds for  $b$  and each  $o(e)$ ).

Let  $o$  be any circuit of  $E$ . It remains to show that  $\sum_{e \in o \cap b} c_o(e) d_b(e) = 0$ . Plugging in the values for  $d_b(e)$ , it remains to show

$$\hat{c}_o(e_b) - \sum_{e \in o \cap (b - e_b)} \frac{c_o(e) c_o(e)(e_b)}{c_o(e)(e)} = 0.$$

That is, we need  $c(e_b) = 0$ , where

$$c = \hat{c}_o - \sum_{e \in o \cap (b - e_b)} \frac{c_o(e)}{c_o(e)(e)} \hat{c}_o(e).$$

As  $c$  is a finite sum of linear dependences, it is again a linear dependence. But for any  $e \in b - e_b$ , we have  $c(e) = \hat{c}_o(e) - \frac{\hat{c}_o(e)}{c_o(e)(e)} c_o(e)(e) = 0$ . By Lemma 2.11, the support of  $c$  is a scrawl, so it can't meet  $b$  in only the point  $e_b$ . Thus  $c(e_b) = 0$ , as desired.  $\square$

Key to this proof is the idea that the painting of the cocircuits of  $M$  is determined by that of the circuits and vice versa. This fact also allows us to observe that a painting of  $M$  uniquely induces paintings of all minors of  $M$ .

**Definition 4.3.** Let  $M$  be a matroid, and let  $((c_o | o \in \mathcal{C}(M)), (d_b | b \in \mathcal{C}(M^*)))$  be a  $k$ -painting of  $M$ . Let the ground set  $E$  of  $M$  be partitioned as  $X \dot{\cup} C \dot{\cup} D$ , and let  $M' = M/C \setminus D$ . We say that a painting  $((c'_o | o \in \mathcal{C}(M')), (d'_b | b \in \mathcal{C}(M'^*)))$  of  $M'$  is *induced* by that of  $M$  if and only if for each  $o' \in \mathcal{C}(M')$  there is  $o \in \mathcal{C}(M)$  with  $o' \subseteq o \subseteq o' \cup C$  and such that  $c'_o = c_o \upharpoonright_{o'}$ , and for each  $b' \in \mathcal{C}(M'^*)$  there is  $b \in \mathcal{C}(M^*)$  with  $b' \subseteq b \subseteq b' \cup D$  and such that  $d'_{b'} = d_b \upharpoonright_{b'}$ .

It is clear from Lemma 2.3 that any painting of  $M$  induces at least one painting of each minor  $M'$ . We can use the fact that the paintings of the circuits and cocircuits determine each other to show that these induced paintings are unique up to scalar factors on the  $c'_o$  and  $d'_b$ .

**Lemma 4.4.** *Let  $M, M'$  and their paintings be as in Definition 4.3. Let  $o'$  be any circuit of  $M'$  and  $o$  any circuit of  $M$  with  $o' \subseteq o \subseteq o' \cup C$ . Then there is  $\lambda \in k^*$  with  $c_o \upharpoonright_{o'} = \lambda c'_{o'}$ .*

*Proof.* Pick any  $e \in o'$ , and let  $\lambda = \frac{c_o(e)}{c'_{o'}(e)}$ . For any other  $f \in o'$ , by Lemma 2.2 there is a cocircuit  $b'$  of  $M'$  with  $o' \cap b' = \{e, f\}$ . Since the painting of  $M'$  is induced from that of  $M$ , there is a cocircuit  $b$  of  $M$  such that  $d_b(g) = d'_{b'}(g)$  for

all  $g \in E(M')$  and  $b' \subseteq b \subseteq b' \cup D$ , and so  $o \cap b = \{e, f\}$ . Using the identities in the definition of painting, we deduce that

$$c_o(e)d_b(e) + c_o(f)d_b(f) = 0 \quad \text{and} \quad c'_{o'}(e)d'_{b'}(e) + c'_{o'}(f)d'_{b'}(f) = 0$$

and so

$$c_o(f) = -\frac{c_o(e)d_b(e)}{d_b(f)} = -\frac{(\lambda c'_{o'}(e))d'_{b'}(e)}{d'_{b'}(f)} = \lambda c'_{o'}(f)$$

which gives the desired result, since  $f$  was arbitrary.  $\square$

Our main result is the following.

**Theorem 4.5.** *Let  $M$  be a tame matroid and  $k$  be a finite field. Then the following are equivalent.*

1.  $M$  is a thin sums matroid over  $k$ .
2.  $M$  is  $k$ -paintable.
3. Every finite minor of  $M$  is  $k$ -representable.

In Theorem 3.1 we already proved this theorem if  $k = \mathbb{F}_2$ . The general case uses similar ideas but is more complex.

*Proof.* To see that (1) implies (3), we use that the class of tame thin sums matroids is closed under taking minors, as shown in [1].

To see that (2) implies (1), let  $((c_o|o \in \mathcal{C}(M)), (d_b|b \in \mathcal{C}(M^*)))$  be a  $k$ -painting. From this  $c$  we can construct a representation of  $M$  as a thin sums matroid over  $k$ . Let  $E \xrightarrow{f} k^{\mathcal{C}(M^*)}$  be the function sending  $e$  to the function sending  $b$  to  $d_b(e)$  if  $e \in b$  and to 0 otherwise. For any thin dependence  $c$  and any cocircuit  $b$  we know that  $0 = \sum_{e \in E} c(e)f(e)(b) = \sum_{e \in b} c(e)d_b(e)$ . In particular, the support of  $c$  never meets a cocircuit only once, so it is a scrawl by Lemma 2.6 and so  $M$ -dependent if it is nonempty. Thus every  $M_{ts}(f)$ -dependent set is  $M$ -dependent.

Conversely, for any  $M$ -circuit  $o$  the function from  $E$  to  $k$  sending  $e$  to  $c_o(e)$  if  $e \in o$  and to 0 otherwise is a thin dependence for  $f$  with support equal to that circuit, and so any  $M$ -dependent set is also  $M_{ts}(f)$ -dependent.

It remains to show that (3) implies (2). We will use a compactness argument, so we begin by defining the topological space we will use. We would like an element of of this space to correspond to a choice of functions as in Lemma 4.1 (though not necessarily satisfying the restrictions of that Lemma), so we take

$$H = \left( \bigcup_{o \in \mathcal{C}(M)} \{o\} \times o \right) \amalg \left( \bigcup_{b \in \mathcal{C}(M^*)} \{b\} \times b \right)$$

and take the underlying set of our space to be  $X = (k^*)^H$  - the compact topology on  $X$  that we will use is given by the product of  $H$  copies of the discrete topology on  $k^*$ .

For each circuit  $o$  and cocircuit  $b$  of  $M$ , the set

$$C_{o,b} = \left\{ c \in (k^*)^H \mid \sum_{e \in o \cap b} c(o, e)c(b, e) = 0 \right\}$$

is closed because  $o \cap b$  is finite. We shall now show that any finite intersection of such sets is nonempty.

Let  $K \subseteq \mathcal{C}(M) \times \mathcal{C}(M^*)$  be finite. Let  $O$  be the set of circuits appearing as first components of elements of  $K$ , and let  $B$  be the set of cocircuits appearing as second components of elements of  $K$ . Let  $F = \bigcup O \cap \bigcup B$ . Note that  $F$  is finite, since  $M$  is tame.

Next, we shall construct a finite minor  $M'$  of  $M$  that will help us to prove that the finite intersection is nonempty.

**Lemma 4.6.** *There exists a finite minor  $M'$  of  $M$  such that for every  $o \in O$  there is an  $M'$ -circuit  $o' \subseteq o$  such that  $o' \cap F = o \cap F$  and for every  $b \in B$  there is an  $M'$ -cocircuit  $b' \subseteq b$  such that  $b' \cap F = b \cap F$ .*

*Proof of the lemma.* We may assume that for each  $o \in O$  and  $b \in B$  the sets  $o \cap F$  and  $b \cap F$  are nonempty by adding an edge from  $o$  or  $b$  to  $F$  if necessary. We pick an element  $e_o \in o \cap F$  for each  $o \in O$ . Next, for each  $o \in O$  and each  $e \in o \cap F - e_o$  we pick a cocircuit  $b_{o,e}$  with  $o \cap b_{o,e} = \{e_o, e\}$  (this is possible by Lemma 2.2). Let  $B'$  be the set of all cocircuits picked in this way or contained in  $B$ . Note that  $B'$  is finite. Similarly, we pick for each  $b \in B$  an element  $e_b \in F \cap b$  and then pick a circuit  $o_{b,e}$  with  $o_{b,e} \cap b = \{e_b, e\}$  for each  $e \in F \cap b - e_b$ , and we collect all of these, together with all circuits contained in  $O$ , in a finite set  $O'$ .

Let  $F' = \bigcup O' \cap \bigcup B'$ . Note that  $F'$  is also finite since  $M$  is tame. Let  $C = \bigcup O' \setminus F'$ , and let  $D = E \setminus \bigcup O'$ . Thus  $E = C \dot{\cup} F' \dot{\cup} D$ . Let  $M'$  be the finite minor of  $M$  with ground set  $F'$  that is given by  $M/C \setminus D$ . For each  $o \in O$ ,  $o \setminus F' \subseteq C$  and so  $o \cap F'$  is a scrawl of  $M'$  by Corollary 2.7. Let  $o'$  be a circuit of  $M'$  with  $e_o \in o' \subseteq o \cap F'$ . Then for each  $e \in o \cap F - e_o$  we know that  $F' \cap b_{o,e}$  is a coscawl of  $M'$ , again by Corollary 2.7, so it can't meet  $o'$  in just one point. But  $e_o \in o' \cap F' \cap b_{o,e} \subseteq \{e_o, e\}$  so we must have  $o' \cap F' \cap b_{o,e} = \{e_o, e\}$  and we conclude that  $e \in o'$ . Since  $e$  was arbitrary, this implies that  $o \cap F \subseteq o'$ . Moreover,  $o \cap F = o' \cap F$ .

Similarly, for each  $b \in B$ , we find a cocircuit  $b'$  of  $M'$  such that  $e_b \in b' \subseteq F' \cap b$ , and it follows by a dual argument that  $F \cap b = b' \cap F$ , completing the proof of the lemma.  $\square$

Since  $M'$  is finite, it is representable over  $k$ . So we can find functions  $c_o$  and  $d_b$  as in Lemma 4.2 for this matroid. Let  $c \in (k^*)^H$  be chosen so that, for each  $o \in O$  and each  $e \in o \cap F$  we have  $c(o, e) = c_{o'}(e)$ , and also so that for each  $b \in B$  and each  $e \in F \cap b$  we have  $c(b, e) = c_{b'}(e)$ . These choices ensure that  $c \in \bigcap_{(o,b) \in K} C_{o,b}$ .

Since  $(k^*)^H$  is compact, and any finite intersection of the  $C_{o,b}$  is nonempty, we have that  $\bigcap_{(o,b) \in \mathcal{C}(M) \times \mathcal{C}(M^*)} C_{o,b}$  is nonempty. As any element in the intersection is a  $k$ -painting, this completes the proof.  $\square$

We note that this gives a uniform way to extend excluded minor characterizations of representability from finite to infinite matroids. For example, we may immediately extend the result of [2, 11] as follows:

**Corollary 4.7.** *A tame matroid  $M$  is a thin sums matroid over  $GF(3)$  if and only if it has no minor isomorphic to  $U_{2,5}$ ,  $U_{3,5}$ ,  $F_7$  or  $F_7^*$ .*

## 5 Other applications of the method

### 5.1 Regular matroids

A key definition to prove Theorem 4.5 was that of a  $k$ -painting. The corresponding notion for regular matroids is as follows.

A *signing* for a matroid  $M$  is a choice of functions  $c_o: o \rightarrow \{1, -1\}$  for each circuit  $o$  of  $M$  and for each cocircuit  $b$  of  $M$  a function  $d_b: b \rightarrow \{1, -1\}$  such that for any circuit  $o$  and cocircuit  $b$  we have

$$\sum_{e \in o \cap b} c_o(e) d_b(e) = 0,$$

where the sum is evaluated over  $\mathbb{Z}$ . A matroid is *signable* if it has a signing.

**Lemma 5.1.** *[[10, Proposition 13.4.5]] Let  $M$  be a finite matroid. Then  $M$  is regular if and only if  $M$  is signable.*

Using similar ideas to those in the proof of Theorem 4.5, we obtain the following.

**Theorem 5.2.** *Let  $M$  be a tame matroid. Then the following are equivalent.*

1.  $M$  is a thin sums matroid over every field.
2.  $M$  is signable
3. Every finite minor of  $M$  is regular.

*Proof.* (2) implies that  $M$  is  $k$ -paintable for every field  $k$ , and so implies (1). (1) implies that every finite minor of  $M$  is representable over every field, and so is regular, which gives (3). (3) implies that every finite minor of  $M$  is signable, by Lemma 5.1. We may then deduce (2) by a compactness argument like that in the proof of Theorem 4.5.  $\square$

Motivated by this theorem, we call a tame matroid *regular* if any of these equivalent conditions hold.

## 5.2 Partial fields

Theorem 5.2 is a special case of a more general result extending characterisations of simultaneous representations over multiple fields using partial fields to tame infinite matroids. For some background on partial fields, see [13].

A *partial field* consists of a pair  $(R, S)$ , where  $R$  is a ring and  $S$  is a subgroup of the group of units of  $R$  under multiplication, such that  $-1 \in S$ . In this context, an  $(R, S)$ -*painting* for a matroid  $M$  is a choice of functions  $c_o: o \rightarrow S$  for each circuit  $o$  of  $M$  and for each cocircuit  $b$  of  $M$  a function  $d_b: b \rightarrow S$  such that for any circuit  $o$  and cocircuit  $b$  we have

$$\sum_{e \in o \cap b} c_o(e) d_b(e) = 0. \quad (2)$$

For example, for any field  $k$  a matroid  $M$  is  $k$ -paintable if and only if it is  $(k, k^*)$ -paintable, and  $M$  is signable if and only if it is  $(\mathbb{Z}, \{-1, 1\})$ -paintable. It is clear that the class of  $(R, S)$ -paintable matroids is closed under duality and under taking minors. In particular, any finite minor of an  $(R, S)$ -paintable matroid is  $(R, S)$ -paintable. The converse follows from an almost identical compactness argument to that used for Theorem 4.5, giving:

**Theorem 5.3.** *Let  $(R, S)$  be a partial field with  $S$  finite. A tame matroid is  $(R, S)$ -paintable if and only if all its finite minors are.*

It follows from the results of [13, Section 2.7] that a finite matroid is  $(R, S)$ -paintable if and only if it is  $(R, S)$ -representable. For finite matroids it is known that simultaneous representability over sets of fields corresponds to representability over partial fields, and we are now in a position to lift many such results to all tame matroids. For example, we can lift [14, Theorem 1.2] as follows:

**Corollary 5.4.** *A tame matroid  $M$  is a thin sums matroid over both  $\mathbb{F}_3$  and  $\mathbb{F}_4$  if and only if it is  $(\mathbb{C}, \{\zeta^i | i \leq 6\})$ -paintable for  $\zeta$  a primitive sixth root of unity.*

## 5.3 Ternary matroids

For finite matroids, a useful property of  $\mathbb{F}_3$ -representable matroids is the uniqueness of the representations. In this section, we shall prove the corresponding property for tame ternary matroids.

Let  $M$  be a  $k$ -paintable matroid for some field  $k$ . We say that two  $k$ -paintings  $((c_o | o \in \mathcal{C}(M)), (d_b | b \in \mathcal{C}(M^*)))$  and  $((\tilde{c}_o | o \in \mathcal{C}(M)), (\tilde{d}_b | b \in \mathcal{C}(M^*)))$  are *equivalent* if and only if there are constants  $x(o)$  for every  $o \in \mathcal{C}(M)$ , constants  $x(b)$  for every  $b \in \mathcal{C}(M^*)$ , constants  $x(e)$  for every edge  $e$  and a field automorphism  $\varphi$  such that the following are true:

1.  $\tilde{c}_o(e) = \varphi(x(o)x(e)c_o(e))$  for any  $e \in o \in \mathcal{C}(M)$ .
2.  $\tilde{d}_b(e) = \varphi\left(\frac{x(b)d_b(e)}{x(e)}\right)$  for any  $e \in b \in \mathcal{C}(M^*)$ .

Two signings of the same matroid  $M$  are *equivalent* if and only if they induce equivalent  $\mathbb{F}_3$ -paintings of  $M$ .

Via Lemma 4.1 for any tame matroid any thin sums representation over  $k$  corresponds to a  $k$ -painting. For finite matroids, the notions of equivalence for representations and paintings coincide: it is straightforward to check that two representations are equivalent iff the corresponding paintings are. As for finite matroids, we obtain the following.

**Theorem 5.5.** *Any two  $\mathbb{F}_3$ -paintings of the same matroid  $M$  are equivalent.*

*Proof.*  $M$ , being  $\mathbb{F}_3$ -paintable, must be tame. Without loss of generality we may also assume that  $M$  is connected and has more than one edge. Thus any edges  $e$  and  $f$  of  $M$  lie on a common circuit<sup>1</sup>. We nominate a particular edge  $g_1$ , and for each other edge  $g$  we nominate a circuit  $o(g)$  containing both  $g_1$  and  $g$ . We also nominate for each circuit  $o$  of  $M$  an edge  $e(o) \in o$  and for each cocircuit  $b$  of  $M$  an edge  $e(b) \in b$ .

We denote the two  $\mathbb{F}_3$ -paintings  $((c_o|o \in \mathcal{C}(M)), (d_b|b \in \mathcal{C}(M^*)))$  and  $((\tilde{c}_o|o \in \mathcal{C}(M)), (\tilde{d}_b|b \in \mathcal{C}(M^*)))$ . We shall construct witnesses to the equivalence as in the definition above. Since every automorphism of  $\mathbb{F}_3$  is trivial, we shall take  $\varphi$  to be the identity.

We now set  $x(g) = \frac{\tilde{c}_o(g)c_o(g_1)}{\tilde{c}_o(g_1)c_o(g)}$  for each  $g \in E$ ,  $x(o) = \frac{\tilde{c}_o(e(o))}{x(e(o))c_o(e(o))}$  for each circuit  $o$  of  $M$  and  $x(b) = \frac{x(e(b))\tilde{d}_b(e(b))}{d_b(e(b))}$  for each cocircuit  $b$  of  $M$ .

In order to prove that these values satisfy (1) at a particular circuit  $o$  and  $g \in o$ , let  $O = \{o, o(g), o(e(o))\}$  and  $F = \{g, g_1, e(o)\}$  and use the construction from the proof of Lemma 4.6 to obtain a finite minor  $M' = M/C \setminus D$  such that for every  $o \in O$  there is an  $M'$ -circuit  $o' \subseteq o$  such that  $o' \cap F = o \cap F$  and for every  $b \in B$  there is an  $M'$ -cocircuit  $b' \subseteq b$  such that  $b' \cap F = b \cap F$ .

Let  $((c'_o|o \in \mathcal{C}(M')), (d'_b|b \in \mathcal{C}(M'^*)))$  be the  $\mathbb{F}_3$ -painting of  $M'$  induced by  $((c_o|o \in \mathcal{C}(M)), (d_b|b \in \mathcal{C}(M^*)))$ , and  $((\tilde{c}'_o|o \in \mathcal{C}(M')), (\tilde{d}'_b|b \in \mathcal{C}(M'^*)))$  that induced by  $((\tilde{c}_o|o \in \mathcal{C}(M)), (\tilde{d}_b|b \in \mathcal{C}(M^*)))$ .

By uniqueness of representation for finite matroids, we can find constants  $x'(o')$  for every  $o' \in \mathcal{C}(M')$ , constants  $x'(b')$  for every  $b' \in \mathcal{C}(M'^*)$  and constants  $x'(g)$  for every  $g \in X$  such that

3.  $\tilde{c}'_{o'}(g) = x'(o')x'(g)c'_{o'}(g)$  for any  $g \in o' \in \mathcal{C}(M')$ .
4.  $\tilde{d}'_{b'}(g) = \frac{x'(b')d'_{b'}(g)}{x'(g)}$  for any  $g \in b' \in \mathcal{C}(M'^*)$ .

**Lemma 5.6.** *For each  $o \in O$  there is  $\lambda_o \in k^*$  such that*

5.  $c_o \upharpoonright_F = \lambda_o c'_{o'} \upharpoonright_F$

*Proof.* As part of the construction of  $M'$ , we picked a canonical element  $e_o$  of  $o'$ . Let  $\lambda = \frac{c_o(e_o)}{c'_{o'}(e_o)}$ . For any other  $e \in o' \cap F$ , there is by construction a cocircuit

<sup>1</sup>In Section 3 of [6], it is shown that the relation ‘ $e$  is in a common circuit with  $f$ ’ is indeed an equivalence relation for infinite matroids.



$b_{o,e}$  of  $M$  with  $o \cap b_{o,e} = \{e_o, e\}$ . Then by the dual of Corollary 2.7  $b_{o,e} \cap E(M')$  is a coscrawl of  $M'$ , and so there is a cocircuit  $b'$  of  $M'$  with  $e_o \in b' \subseteq b_{o,e}$ , and so  $e_o \in o' \cap b' \subseteq \{e_o, e\}$ . Since  $o'$  and  $b'$  can't meet in only one element,  $e \in b'$ . Since the painting of  $M'$  is induced from that of  $M$ , there is a cocircuit  $b$  of  $M$  such that  $d_b(e) = d'_{b'}(e)$  for all  $e \in E(M')$  and  $b' \subseteq b \subseteq b' \cup D$ , and so  $o \cap b = \{e_o, e\}$ . Using the identities in the definition of painting, we deduce that

$$c_o(e_o)d_b(e_o) + c_o(e)d_b(e) = 0 \quad \text{and} \quad c'_{o'}(e_o)d'_{b'}(e_o) + c'_{o'}(e)d'_{b'}(e) = 0$$

and so

$$c_o(e) = -\frac{c_o(e_o)d_b(e_o)}{d_b(e)} = -\frac{\lambda c'_{o'}(e_o)d'_{b'}(e_o)}{d'_{b'}(e)} = \lambda c'_{o'}(e)$$

which gives the desired result, since  $e \in o' \cap F$  was arbitrary.  $\square$

Similarly, we can find constants  $\tilde{\lambda}_o$  for each  $o \in O$  such that

$$6. \quad \tilde{c}_o \upharpoonright_F = \tilde{\lambda}_o \tilde{c}'_{o'} \upharpoonright_F$$

Now we must simply unwind all the algebraic relationships to obtain the desired result.

$$x(g) = \frac{\tilde{c}_{o(g)}(g)c_{o(g)}(g_1)}{\tilde{c}_{o(g)}(g_1)c_{o(g)}(g)} = \frac{\tilde{c}'_{o(g)'}(g)c'_{o(g)'}(g_1)}{\tilde{c}'_{o(g)'}(g_1)c'_{o(g)'}(g)} = \frac{x'(o(g)')x'(g)}{x'(o(g)')x'(g_1)} = \frac{x'(g)}{x'(g_1)}$$

where the first equation follows from the definitions, the second from (5) and (6) and the third from (3). Similarly, we get:

$$x(o) = \frac{\tilde{c}_o(e(o))}{x(e(o))c_o(e(o))} = \frac{\tilde{\lambda}_o}{\lambda_o} \frac{\tilde{c}'_{o'}(e(o))}{x(e(o))c'_{o'}(e(o))} = \frac{\tilde{\lambda}_o}{\lambda_o} \frac{x'(o')x'(e(o))}{x(e(o))}$$

And finally:

$$x(o)x(g)c_o(g) = \frac{\tilde{\lambda}_o}{\lambda_o} \frac{x'(o')x'(e(o))x'(g_1)}{x'(e(o))} \frac{x'(g)}{x'(g_1)} c_o(g) = \frac{\tilde{\lambda}_o}{\lambda_o} x'(o')x'(g)c_o(g)$$

Now the last term is just  $\tilde{c}_o(g)$  by first applying (5) and then (3). This completes the proof of the above assignment satisfies (1). The proof that it also satisfies (2) is similar.  $\square$

As every tame regular matroid is a thin sums matroid over  $\mathbb{F}_3$ , it also has a unique representation. In particular the finite cycle matroid, the algebraic cycle matroid and the topological cycle matroid of a given graph (and their duals) have a unique signing.

In what follows, we will describe this signing of the finite cycle matroid of a given graph  $G$ . First direct the edges of  $G$  in an arbitrary way. To define the

functions  $c_o$ , let  $o$  be some cycle of  $G$ . Pick a cyclic order of  $o$ . For  $e \in o$ , let  $c_o(e) = 1$  if  $e$  is directed according to the cyclic order of  $o$  and  $-1$  else.

Next, let  $b$  be some cocircuit. By minimality of the cocircuit, it is contained in a single component of  $G$  and its removal separates this component into two components, say  $C_1(b)$  and  $C_2(b)$ . Note that every edge in  $b$  has precisely one endvertex in each of these components. For  $e \in b$ , let  $d_b(e) = 1$  if  $e$  points to a vertex in  $C_1$  and  $-1$  else.

It is worth noting that the directions we gave to the edges of  $G$  do not influence the values of the products  $c_o \cdot d_b$ .

It remains to check that  $\sum_{e \in o \cap b} c_o(e)d_b(e) = 0$  for all circuits  $o$  and cocircuits  $b$ . As every circuit is finite, the above sum is finite. To see that the above formula is true, it suffices to show  $c_o(e)d_b(e) = -c_o(f)d_b(f)$  for any two edges  $e, f \in b \cap o$  that can be joined by a subpath  $P(e, f)$  of  $o$  only containing edges outside  $b \cap o$ . Indeed, then  $o \cap b$  contains as many edges that contribute 1 to the above sum as it contains edges that contribute  $-1$ .

As the value of  $c_o \cdot d_b$  does not depend on the chosen directions of the edges of  $G$ , we may assume that all edges of  $G$  are directed from  $C_1(b)$  to  $C_2(b)$ . First we consider the case when  $c_o(e)d_b(e) = 1$ . Then  $o$  traverses the cocircuit through  $e$  from  $C_2(b)$  to  $C_1(b)$ . As  $P(e, f)$  is completely contained in  $C_1(b)$ , along the edge  $f$  the cycle  $o$  has to traverse the cocircuit from  $C_1(b)$  to  $C_2(b)$ . Thus  $c_o(f)d_b(f) = (-1) \cdot 1 = -1$ , as desired. The case where  $c_o(e)d_b(e) = -1$  is analogue. This completes the construction of the signing of the finite cycle matroid of  $G$ .

Let us look at how to modify the above construction to make it work for the algebraic cycle matroid and the topological cycle matroid instead. It is easy to show that in the 1-point compactification of a locally finite graph, the closure of the edge set of any circuit is homeomorphic to the unit circle. So we also get a cyclic order on the circuits of the algebraic cycle matroid and the above argument applies as the algebraic cycle matroid is tame.

If  $G$  is not locally finite, then  $G$  considered as a topological space is not locally compact, and so has no 1-point compactification - but still a similar construction shows the assertion in this case. A similar construction works for the topological cycle matroid.

## References

- [1] H. Afzali and N. Bowler. Thin sums matroids and duality. In preparation.
- [2] Robert E Bixby. On reid's characterization of the ternary matroids. *Journal of Combinatorial Theory, Series B*, 26(2):174 – 204, 1979.
- [3] N. Bowler and J. Carmesin. Matroids with an infinite circuit-cocircuit intersection. Submitted 2012, current version available at <http://www.math.uni-hamburg.de/spag/dm/papers/intersection.pdf>.

- [4] N. Bowler, J. Carmesin, and R. Christian. Graph-like spaces and infinite matroids. In preparation.
- [5] H. Bruhn and R. Diestel. Infinite matroids in graphs. arXiv:1011.4749 [math.CO], 2010.
- [6] H. Bruhn, R. Diestel, M. Kriesell, R. Pendavingh, and P. Wollan. Axioms for infinite matroids. arXiv:1003.3919 [math.CO], 2010.
- [7] Henning Bruhn and Paul Wollan. Finite connectivity in infinite matroids. *European Journal of Combinatorics*, 33(8):1900 – 1912, 2012.
- [8] R. Diestel. *Graph Theory* (4th edition). Springer-Verlag, 2010. Electronic edition available at: <http://diestel-graph-theory.com/index.html>.
- [9] J. Geelen, A. Gerards, and A. Kapoor. The excluded minors for GF(4)-representable matroids. *J. Combin. Theory (Series B)*, 79:247–299, 2000.
- [10] J. Oxley. *Matroid Theory*. Oxford University Press, 1992.
- [11] P.D Seymour. Matroid representation over  $\text{gf}(3)$ . *Journal of Combinatorial Theory, Series B*, 26(2):159 – 173, 1979.
- [12] W. T. Tutte. A homotopy theorem for matroids, ii. *Transactions of the American Mathematical Society*, 88(1):pp. 161–174, 1958.
- [13] S.H.M. van Zwam. *Partial fields in matroid theory*. PhD thesis, Eindhoven University of Technology, 1999.
- [14] Geoff Whittle. On matroids representable over  $\text{gf}(3)$  and other fields. *Transactions of the American Mathematical Society*, 349(2):pp. 579–603, 1997.