

Analysis III
for Engineering Students
Homework sheet 6

Exercise 1:

a) For the function

$$f : Q \rightarrow \mathbb{R}, \quad f(x, y) = 6 - 2x + 4y$$

with $Q := [0, 3] \times [0, 2]$ compute

(i) Riemannian upper and lower sum for the following equidistant decomposition Z of Q

$$Q_{i,j} = [x_{i-1}, x_i] \times [y_{j-1}, y_j], \quad i, j = 1, \dots, n$$

$$\text{where } x_i = \frac{3i}{n} \text{ and } y_j = \frac{2j}{n}$$

(ii) and the integral of f over Q using Fubini's theorem.

b) (i) Draw the area P enclosed by the functions $f(x) = 2x$ and $g(x) = 24 - 2x^2$ and represent it as the “normal” area.

(ii) Compute $\int_P x \, d(x, y)$.

Exercise 2:

Draw the half cylinder Z given by $1 \leq z \leq 2$, $0 \leq y$ and $x^2 + y^2 \leq 9$ and calculate its center of mass with the density function $\rho(x, y, z) = z$ using cylindrical coordinates.

Exercise 3:

- a) For the vector field $\mathbf{f} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ with $\mathbf{f}(x, y) = \begin{pmatrix} y + \sin x \\ xy^2 \end{pmatrix}$ calculate the integral of the curve (line integral) $\oint_{\mathbf{c}} \mathbf{f}(\mathbf{x}) d\mathbf{x}$.

Here $\mathbf{c}$ is the mathematically positive boundary curve of the area G enclosed by $x^2 \leq y \leq x$ with $0 \leq x \leq 1$.

- b) For the vector field $\mathbf{f} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ with $\mathbf{f}(x, y, z) = \begin{pmatrix} -z^2/2 \\ 0 \\ xz \end{pmatrix}$

calculate the line integral $\int_{\mathbf{c}} \mathbf{f}(\mathbf{x}) d\mathbf{x}$ with the line

$$\mathbf{c} : \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] \rightarrow \mathbb{R}^3 \quad \text{and} \quad \mathbf{c}(t) = \begin{pmatrix} 2 \cos^2 t \\ 2 \sin t \cos t \\ 2 \sin t \end{pmatrix}.$$

Submission deadline: 20.1.2023