

## Differential Equations II for Engineering Students

### Homework sheet 3

#### Exercise 1: [4 +2 Points]

a) Given the following initial-value problem for  $u(x, t)$ ,  $u : \mathbb{R} \times \mathbb{R}^+ \rightarrow \mathbb{R}$

$$\begin{aligned}u_t + u \cdot u_x &= 0, & x \in \mathbb{R}, t \in \mathbb{R}^+ \\u(x, 0) &= g(x), & x \in \mathbb{R}.\end{aligned}$$

Here let  $g : \mathbb{R} \rightarrow \mathbb{R}$  be a strictly monotonically increasing function with two points of discontinuity (jump points).

For each of the following statements, determine if it is true or false.

- (i) There is a unique weak solution.
- (ii) In order to obtain the entropy solution, one has to introduce two shock waves.
- (iii) The entropy solution is valid for all times, i.e. for any  $t \in \mathbb{R}^+$ .

**Justify your answers.**

b) What is the jump condition for the weak solution to

$$\begin{aligned}u_t + (u^3)_x &= 0, & x \in \mathbb{R}, t \in \mathbb{R}^+ \\u(x, 0) &= \begin{cases} 4 & \text{for } x \leq 0, \\ 2 & \text{for } x > 0? \end{cases}\end{aligned}$$

**Exercise 2:** Determine the entropy solution to the Burgers' equation  $u_t + uu_x = 0$  with the initial data

$$u(x, 0) = \begin{cases} 0 & x < 0 \\ 1 & 0 \leq x \leq 1 \\ 0 & x > 1 \end{cases}$$

at the time  $t = 2$ . What new problem occurs at  $t = 2$ ?

**Alternatively:** Determine the solution for  $t > 2$ .

**Exercise 3:**

We discuss again the simple traffic flow model from Sheet 1 with the notation introduced there:

$u(x, t)$  = density of vehicles (vehicles/length) at point  $x$  at time  $t$ ,

$v(x, t)$  = velocity at point  $x$  at time  $t$ ,

$q(x, t) = u(x, t) \cdot v(x, t)$  = flow = number of vehicles passing  $x$  at time  $t$  per time unit.

We improve our model from Sheet 2 by incorporating maximal density and a maximal velocity

$u_{max}$  = maximal density of vehicles (bumper to bumper),

$v_{max}$  = maximal velocity

This can be done, for example, as follows:

$$v(x, t) := v(u(x, t)) := v_{max} \left( 1 - \frac{u(x, t)}{u_{max}} \right)$$

- Set up the continuity equation ( $u_t + q_x = 0$ ).
- Show again that the characteristics are straight lines and determine their slopes.
- Sketch the characteristics for

$$v_{max} = 1 \quad (\text{Here has been scaled appropriately!})$$

$$u(x, 0) = \begin{cases} u_l = u_{max}/2 & x < 0 \\ u_r = u_{max} & x > 0 \end{cases} \quad (\text{red traffic light/traffic jam etc.})$$

- For the Burgers' equation we allowed shock waves only in the case  $u_l > u_r$ . There must obviously be a different condition here. What could be the reason for that?

**Note:** This question can not be answered completely only with help of the lecture slides. You can only make a guess here!

**Abgabetermine: 08.05.-12.05.2023**